

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1966

[Full Table of Content](#)



66/R/CC/C/29 and 66/W/WCD/C/29 Dd and Dazomet - Spring Wheat

Rothamsted Research

Rothamsted Research (1967) *66/R/CC/C/29 and 66/W/WCD/C/29 Dd and Dazomet - Spring Wheat ; Yields Of The Field Experiments 1966*, pp 225 - 226 - DOI:
<https://doi.org/10.23637/ERADOC-1-158>

66/c/29.1

DD AND DAZOMET - SPRING WHEAT

(CC and WCD)

Effects of soil fumigants on yield and soil-borne pathogens, Rothamsted (R) Hoosfield and Woburn (W) Lansome, the first year.

Design: 3 randomised blocks of 3 plots, split into 8.

Area of each sub plot: 0.0048. Area harvested:
Hoosfield (R) - 0.0048, Lansome (W) - 0.0036.

Treatments: All combinations of:-

Whole plots: 1. Nitrogen: 0.5 (N1), 1.0 (N2), 1.5 cwt N (N3)
applied as 'Nitro-Chalk'.

Sub plots: 2. Fumigants: None (0), none, rotary cultivated (R).
DD: 200 (D2), 400 (D4), 800 lb (D8) injected
to 6 inch depth.
Dazomet: 100 (Z1), 200 (Z2), 400 lb (Z4) rotary
cultivated in.

Basal applications: 280 lb (0:20:20) combine drilled. Weedkiller
(Lansome (W) only): Ioxynil/mecoprop (Atril C at 5 pints in
35 gals).

Cultivations, etc.:

Hoosfield (R): Ploughed: Oct 28, 1965. DD applied: Nov 8.
Dazomet applied, Z and R plots rotary cultivated: Mar 7, 1966.
Seed drilled at 180 lb: Apr 26. 'Nitro-Chalk' applied:
Apr 27. Combine harvested: Sept 9. Variety: Klokka.
Previous crops: Barley 1964, winter barley 1965.

Lansome (W): Ploughed: Sept 30, 1965. DD applied: Nov 22.
Dazomet applied, Z and R plots rotary cultivated:
Feb 14, 1966. Seed drilled at 170 lb: Apr 13.
'Nitro-Chalk' applied: Apr 25. Weedkiller applied: May 13.
Combine harvested: Sept 7. Variety: Klokka. Previous crops:
Winter wheat 1964, barley 1965.

NOTE: Soil samples were taken for microflora investigations and
plant samples for root-rotting diseases. Samples were also taken
for nematode counts and plant weights, also soil samples for final
nematode population.

Standard errors per plot. Grain.

Hoosfield (R).	Whole plot: 1.19 or 3.9% (4 d.f.)
	Sub plot: 1.73 or 5.6% (42 d.f.)
Lansome (W).	Whole plot: 2.99 or 7.4% (4 d.f.)
	Sub plot: 2.92 or 7.2% (38 d.f.)

66/C/29.2

SUMMARY OF RESULTS

GRAIN

	O	R	D2	D4	D8	Z1	Z2	Z4	Mean
Hoosfield (R)									
	(1) and (2)								(± 0.69)
N1	25.8	26.3	27.1	24.6	25.3	29.0	30.4	32.6	27.6
N2	28.6	33.7	31.0	30.4	27.1	31.2	34.6	32.3	31.1
N3	32.7	35.1	34.4	32.5	33.7	33.2	33.2	35.9	33.9
Mean (± 0.58)	29.0	31.7	30.8	29.1	28.7	31.1	32.7	33.6	30.9

Mean D.M. %: 83.1

Lansome (W)

	(1) and (2)								(± 1.73)
N1	30.5	31.5	35.1	34.2	40.1	32.0	39.7	41.9	35.6
N2	39.2	40.8	41.3	41.6	43.4	43.4	46.2	45.3	42.7
N3	40.1	38.5	43.7	42.4	41.3	43.7	46.4	45.8	42.8
Mean (± 0.97)	36.6	36.9	40.1	39.4	41.6	39.7	44.1	44.3	40.3

Mean D.M. %: 82.8

	Hoosfield (R)	Lansome (W)	
(1)	(± 1.16)	(± 2.34)	For use in vertical and diagonal comparisons
(2)	(± 1.00)	(± 1.69)	For use in horizontal and interaction comparisons