


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1966

[Full Table of Content](#)



66/S/SC/C/30 Intensive Wheat - Winter Wheat

Rothamsted Research

Rothamsted Research (1967) *66/S/SC/C/30 Intensive Wheat - Winter Wheat ; Yields Of The Field Experiments 1966*, pp 227 - 228 - DOI: <https://doi.org/10.23637/ERADOC-1-158>

66/c/30.1

INTENSIVE WHEAT

(SC)

Saxmundham, Oldershaw's and Garner's plots 1966, the first year.

Design: 4 randomised blocks of 5 plots, with plots (excluding leys and beans) split into 3 for N.

Area of each sub-plot: 0.0182. Area harvested: 0.0096.

Treatments: All combinations of:-

1. Crop sequences:

	1966	1967	1968	1969	1970
1	W	W	W	W	W
2	L	W	W	W	W
3	L	Be	W	W	W
4	W	L	Be	W	W
5	W	W	L	Be	W

Where W = Winter wheat, L = Ley - one year Meadow Fescue (cut twice* for hay), Be = Spring beans.

2. Nitrogen to wheat: 0.6 (N1), 1.2 (N2), 1.8 cwt (N3) N per acre as 'Nitro-Chalk'.

* once only in 1966.

Basal application: 500 lb compound (0:20:20). Other applications:
To Meadow Fescue: 1.0 cwt N and 1.0 K2O as muriate of potash in seedbed, 0.5 cwt N top-dressed. N as 'Nitro-Chalk'. Weedkiller to wheat: Dicamba, MCP, mecoprop and TBA (Cambilene at 4 pints in 20 gals).

Cultivations, etc.: Ploughed: Sept 29, 1965. Basal compound fertiliser applied: Sept 30.

Wheat: Seed drilled: Oct 8, 1965. 'Nitro-Chalk' applied: Mar 24, 1966. Weedkiller applied: Apr 29. Combine harvested: Aug 17. Variety: Cappelle.

Grass ley: 'Nitro-Chalk' applied: Mar 24, 1966. Muriate of potash applied, seed drilled at 16 lb: Mar 28. 'Nitro-Chalk' applied: July 13. Cut for hay: Aug 12. Variety: Canadian Meadow Fescue.

NOTES: (1) Yields were taken for winter wheat only.
(2) Estimates of the incidence of take-all (*Ophiobolus graminis*) and eyespot (*Cercospora herpotrichoides*) were made in April and June.

66/c/30.2

Standard error per sub-plot:

Winter wheat, grain: 2.59 or 9.8% (22 d.f.)

SUMMARY OF RESULTS

WINTER WHEAT

GRAIN

N1	N2	N3	Mean
19.4	28.8	31.5	26.6

Mean D.M. %: 83.8