


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1966

[Full Table of Content](#)



66/R/RP9/1/DD/6 Potatoes - Gangrene

Rothamsted Research

Rothamsted Research (1967) *66/R/RP9/1/DD/6 Potatoes - Gangrene* ; Yields Of The Field Experiments 1966, pp 287 - 288 - DOI: <https://doi.org/10.23637/ERADOC-1-158>

66/Da/6.1

POTATOES

(RP 9/1)

Effects of gangrene (*Phoma* spp.) - Little Hoos 1966.

Design: 6 randomised blocks of 2 plots, split into 4.

Area of each plot: 0.0067. Area harvested: 0.0033.

Treatments: All combinations of:-

Whole plots: 1. Varieties: King Edward (E), Majestic (M).

Sub plots: 2. Levels of seed-infection (*Phoma* spp.): Clean (A), moderately infected (B), severely infected (C), unselected stock (D).

Basal applications: 7.75 cwt (17:11:22). Fungicide: Mancozeb at 1.2 lb in 35 gals on 4 occasions.

Cultivations, etc.: Ploughed: Jan 12, 1966. Basal NPK applied: Mar 30. Rotary cultivated, potatoes planted: May 3. Rotary ridged twice: May 28 and June 17. Fungicide applied: June 30, July 23, Aug 8 and 18. Sprayed with undiluted BOV at 21 gals: Sept 16. Haulm destroyed mechanically: Sept 27. Lifted: Sept 29. Previous crops: Barley 1964, fallow 1965.

Standard errors per plot. Total tubers:

Whole plot: 0.708 or 4.0% (5 d.f.)

Sub plot: 0.873 or 4.9% (30 d.f.)

66/Da/6.2

SUMMARY OF RESULTS

	A	B	C	D	Mean
TOTAL TUBERS					
	(1) and (2)				(±0.289)
E	18.71	18.13	17.35	18.44	18.16
M	17.71	18.01	15.36	17.50	17.15
Mean (±0.252)	18.21	18.07	16.36	17.97	17.65
% WARE					
E	93.9	92.0	91.3	92.7	92.5
M	97.6	96.3	94.5	95.5	96.0
Mean	95.8	94.1	92.9	94.1	94.2

(1) (±0.423) For use in vertical and diagonal comparisons

(2) (±0.356) For use in horizontal and interaction comparisons