

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1967

[Full Table of Content](#)



67/R/BJ/C/12 Intensive Winter Wheat - Winter Barley

Rothamsted Research

Rothamsted Research (1968) *67/R/BJ/C/12 Intensive Winter Wheat - Winter Barley* ; Yields Of The Field Experiments 1967, pp 161 - 162 - DOI: <https://doi.org/10.23637/ERADOC-1-157>

67/c/12.1

INTENSIVE WINTER BARLEY GROWING EXPERIMENT

(BJ)

Hoosfield (Old Four Course) 1967 - the third year

For details of treatments, etc. see 'Results' 65/C/25 and for previous years' results see 'Results' 65/C/25 and 66/C/16.

Area of each plot: 0.0386. Area harvested: 0.0254.

Basal applications: Basal fertiliser as previously. 23 cwt ground chalk. Weedkiller to barley and oats: Ioxynil/mecoprop (Atril C at 6 pints in 20 gals). Insecticide to beans: Demeton-s-methyl (Metasystox at 15 fluid oz in 37 gals) by hand on 2 occasions.

Cultivations, etc.: Ground chalk applied: Sept 22, 1966. Ploughed: Oct 17.

Barley: Seed drilled at 140 lb: Oct 28. 'Nitro-Chalk' applied: Apr 7, 1967. Weedkiller applied: May 9. Combine harvested: Aug 7.

Oats: Seed drilled at 145 lb: Oct 29, 1966. 'Nitro-Chalk' applied: Apr 7, 1967. Weedkiller applied: May 9. Combine harvested: Aug 7. Variety: Peniarth.

Winter beans: Seed drilled at 300 lb: Oct 29, 1966. Insecticide applied: June 27 and July 7, 1967. Combine harvested: Sept 9. Variety: Pedigree.

- NOTES: (1) Yields were taken from sequences 1, 3 (Barley) and 4 (Oats) only.
(2) Estimates of Eyespot (*Cercospora herpotrichoides*) and take-all (*Ophiobolus graminis*) were made in spring and summer.

Standard errors per plot. Grain:

Barley Sequences 1 and 3: 2.82 or 11.4% (10 d.f.)
Oats Sequence 4: 1.38 or 4.2% (4 d.f.)

67/C/12.2

SUMMARY OF RESULTS
BARLEY (1 and 3) GRAIN

Crop in 1965	1966	N0	N1	N2	N3	Mean N1 N2
			(±1.63)			(±1.15)
B	B (1)		18.1	27.7	28.6	22.9
B	Be (3)	20.2	28.2	26.2		27.2

General mean: 24.8
Mean D.M. %: 83.9

OATS (4) GRAIN			
N1	N2	N3	Mean
28.3	34.5 (±0.80)	36.5	33.1

Mean D.M. %: 83.0