

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1967

[Full Table of Content](#)



67/R/CA/C/21 Levels and Forms of N for Beans - Winter Wheat

Rothamsted Research

Rothamsted Research (1968) *67/R/CA/C/21 Levels and Forms of N for Beans - Winter Wheat ; Yields Of The Field Experiments 1967*, pp 189 - 190 - DOI: <https://doi.org/10.23637/ERADOC-1-157>

67/C/21.1

LEVELS AND FORMS OF N FOR BEANS

(CA/W)

Levels and forms of N for beans followed by wheat, Great Knott II 1967, the second year - winter wheat.

Design: 3 x 3 x 3 in 3 blocks of 9 plots, plus 3 extra plots per block.

Area of each plot: 0.0193. Area harvested: 0.0129.

Treatments: All combinations of:-

1. Nitrogen to winter wheat 1967: None (N0), 0.5 (N1), 1.0 (N2) cwt N as 'Nitro-Chalk'.

Treatments to beans 1966:-

2. Forms of nitrogen: Ammonium nitrate (A), nitrate of soda (N), sulphate of ammonia (S).
3. Levels of N: None (R0), 1 (R1), 2 (R2), 3 (R3) cwt N broadcast.

Additional plots: To wheat 1967 N0, N1 and N2 as above to plots receiving no nitrogen in 1966.

Basal applications: 340 lb (0:14:28) combine drilled. Weedkiller: Dicamba at 1.28 oz and MCPA at 1.13 lb a.e. in 32 gals.

Cultivations, etc.: Ploughed: Sept 29, 1966. Seed drilled at 180 lb: Oct 29. 'Nitro-Chalk' applied: Apr 17, 1967. Sprayed: Apr 29. Combine harvested: Aug 30. Variety: Cappelle. Previous crops: Barley 1965, spring beans 1966.

NOTE: For the previous year's results see 'Results' 66/C/28.

Standard error per plot.

Grain: 4.24 or 8.0% (9 d.f.)

67/C/21.2

SUMMARY OF RESULTS

GRAIN

	A	N	S	R1	R2	R3	Mean
	(±2.45)			(±2.45)			(±1.41)
NO	47.7	46.4	44.3	38.6	47.1	52.7	46.1
N1	53.4	54.8	57.2	55.4	54.5	55.5	55.1
N2	56.5	57.7	56.1	60.0	57.8	52.5	56.8
				(±2.45)			
	A			53.1	51.4	53.1	52.5
	N			51.5	54.4	52.9	53.0
	S			49.4	53.6	54.7	52.5
Mean (±1.41)				51.3	53.1	53.6	52.7

Plots receiving no nitrogen 1966

NO	N1	N2
40.5	51.9	56.8
	(±2.45)	

General mean: 51.9

Mean D.M. %: 84.9