

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1967

[Full Table of Content](#)



67/W/WOM/C/24 Organic Manuring - Potatoes and Grass

Rothamsted Research

Rothamsted Research (1968) 67/W/WOM/C/24 Organic Manuring - Potatoes and Grass ; Yields Of The Field Experiments 1967, pp 197 - 200 - DOI: <https://doi.org/10.23637/ERADOC-1-157>

67/C/24.1

ORGANIC MANURING EXPERIMENT

(WOM)

The cumulative effects of organic matter on light land - Woburn
Stackyard B 1967, 3rd year.

For previous history, rotation, treatments etc., see 'Results'
66/C/31. All plots except those under leys (LC and LN)
carried potatoes.

Area of each sub plot: 0.0153. Area harvested: Potatoes - 0.0102,
leys - 0.0129.

Fertilisers applied Autumn 1966 (cwt)

Treatment	P2O5	K2O	MgO
DG	-	-	-
ST	0.4	-	0.25
PT	0.5	1.0	-
GM	-	1.0	0.15
FD	2.0	3.0	0.8
FS	0.5	1.0	0.15
LC	0.5	2.0	0.15
LN	0.5	2.0	0.15

Fertilisers applied Spring 1967 (cwt)

Treatment	P2O5	K2O	MgO
DG	-	-	-
ST	1.0	1.0	0.20
PT	1.0	1.0	-
GM	1.0	1.0	0.25
FD	2.5	1.2	0.4
FS	1.0	1.0	0.25
LC	1.0	1.25+1.25*	0.5
LN	1.0	1.5+1.5*	0.5

* After first cut.

Because the peat applied in 1965 and 1966 had less dry matter than ex-
pected additional peat was applied and worked in with the Spring
fertilisers before the potatoes were planted at 6.10 tons fresh
material at 35.5% D.M. equivalent to 2 tons 3 cwt 35 lb dry
matter.

67/C/24.2

Nitrogen to potatoes:

N1, N2, N3, N4. 0.2, 0.6, 1.0, 1.4 cwt N as 'Nitro-Chalk'.

Cultivations, etc.:

LC and LN plots: P,K and Mg applied: Nov 4, 1966. 'Nitro-Chalk' applied to LN plots: Apr 4, 1967, June 19.

P,K and Mg applied: Apr 7. LC plots cut: June 8, July 13.

LN plots cut: June 8.

Potatoes: P,K and Mg applied: Nov 11, 1966. Peat, straw, dung applied: Nov 28 - 29. Ploughed: Nov 30 - Dec 1.

Additional peat applied: Apr 4, 1967. N,P,K and Mg applied: Apr 4 - 6. Rotary cultivated, potatoes planted: Apr 5 - 7.

Earthed up: May 10. Rotary ridged: June 10. Sprayed with mancozeb at 1.2 lb in 30 gals: July 1, July 28, Aug 8.

Sprayed with undiluted BOV at 15 gals: Aug 21. Lifted:

Sept 12 - 13. Variety: Majestic.

Standard errors per plot. Potatoes, total tubers:

Whole plot: 1.340 or 19.2% (15 d.f.)

Sub plot: 0.925 or 13.2% (54 d.f.)

67/C/24.3

SUMMARY OF RESULTS

POTATOES

	NO	N1	N2	N3	Mean
TOTAL TUBERS					
	(1) and (2)				(± 0.670)
DG	5.77	6.93	8.97	10.43	8.02
ST	3.95	5.70	7.90	8.91	6.62
PT	3.61	6.10	5.68	7.83	5.80
GM	6.67	8.28	10.18	11.06	9.05
FD	4.45	5.85	8.25	9.62	7.04
FS	3.67	5.03	6.06	6.97	5.43
Mean (± 0.189)	4.69	6.32	7.84	9.14	6.99

(1) (± 0.780) For use in vertical and diagonal comparisons(2) (± 0.462) For use in horizontal and interaction comparisons

% WARE

DG	85.5	88.9	90.5	89.6	88.6
ST	82.2	83.9	90.3	90.4	86.7
PT	80.0	83.8	87.6	88.2	84.9
GM	85.4	89.6	89.6	91.5	89.0
FD	86.5	87.7	91.4	88.8	88.6
FS	84.4	83.9	87.4	85.4	85.3
Mean	84.0	86.3	89.5	89.0	87.2

67/C/24.4

GRASS: DRY MATTER

	LC	LN
1ST CUT		
	33.1	42.1
2ND CUT		
	2.9	-
TOTAL OF 2 CUTS		
	35.9	42.1
Mean D.M. %: 1st cut:	18.0	24.7
2nd cut:	37.8	-