

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1967

[Full Table of Content](#)



67/R/CU/C/27 Legumes and Barley - Various Crops

Rothamsted Research

Rothamsted Research (1968) *67/R/CU/C/27 Legumes and Barley - Various Crops ; Yields Of The Field Experiments 1967*, pp 207 - 209 - DOI: <https://doi.org/10.23637/ERADOC-1-157>

67/c/27.1

LEGUMES AND BARLEY

(CH)

Effects of crop sequences and green manures - Highfield IV 1967, the first year.

Design: 2 randomised blocks of 12 plots.

Area of each plot: 0.0521. Area harvested: Barley and oats - 0.0276, oilseed rape - 0.0296, red clover - 0.0059, spring beans - 0.0311.

Treatments:

	Crop sequences (C):-						
	1	2	3	4	5	6	7
1967	B	B	O	O	Ra	Cl	Be
Undersown	-	T	-	T	-	-	-
1968	B	B	B	B	B	B	B

With nitrogen in 1967: 0.4 (N1), 0.8 (N2) cwt N as 'Nitro-Chalk' to oats and barley, 0.8 (N2), 1.6 (N4) to oilseed rape, none to beans and clover.

B = Barley, O = Oats, Ra = Oilseed Rape, Cl = Red Clover, Be = Spring beans, T = Trefoil.

Basal and other applications: 280 lb (0:20:20) broadcast, ground chalk at 40 cwt to southern half, at 80 cwt to northern half. Weed-killers: Beans - simazine at 1 lb in 32 gals, barley and oats (excluding undersown plots) - ioxynil/mecoprop (Atril C at 5 pints in 20 gals). Insecticides: Beans - demeton-s-methyl at 3 oz in 37 gals and later at 3.75 oz in 30 gals by hand, oilseed rape - malathion at 18 oz in 30 gals.

Cultivations, etc.: Ploughed: Nov 4, 1966. Ground chalk applied: Jan 6 and Feb 9, 1967. Basal PK compound applied, oats drilled at 160 lb, barley at 140 lb, beans at 200 lb, weedkiller applied to beans: Mar 8. 'Nitro-Chalk' applied to oats and barley: Mar 28. Clover sown at 30 lb: Mar 29. Oilseed rape sown at 7.5 lb: Mar 30. Trefoil sown at 32 lb: May 9. Weedkiller applied to barley and oats: June 1. Insecticides applied: Beans - June 14, rape - June 16, beans - July 7. Clover cut twice: Aug 4 and Oct 2. Combine harvested: Oats - Aug 22, barley - Aug 23, beans - Sept 1, oilseed rape - Sept 6. Varieties: Oats - Manod, barley - Maris Badger, beans - Tarvin, oilseed rape - Nilla, clover - Dorset Marl, trefoil - English (inoculated seed).

67/C/27.2

SUMMARY OF RESULTS

GRAIN

UNDERSOWN

	-	T	Mean
	BARLEY		
N1	31.1	32.3	31.7
N2	36.8	34.0	35.4
Mean	34.0	33.1	33.6

Mean D.M. %: 82.5

	OATS		
N1	32.7	32.3	32.5
N2	36.4	34.5	35.5
Mean	34.6	33.4	34.0

Mean D.M. %: 83.3

67/C/27.3

OILSEED RAPE

SEED

N2	N4	Mean
17.5	16.2	16.9

Mean D.M. %: 85.8

RED CLOVER

	1ST CUT	2ND CUT	TOTAL OF 2 CUTS
Mean	29.2	22.9	52.1
Mean D.M. %:	18.0	14.7	16.4

SPRING BEANS

GRAIN

Mean	30.6
Mean D.M. %:	74.2