

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1967

[Full Table of Content](#)



67/R/EB/C/37 Chemical Control of Take-all - Winter Wheat

Rothamsted Research

Rothamsted Research (1968) *67/R/EB/C/37 Chemical Control of Take-all - Winter Wheat ; Yields Of The Field Experiments 1967*, pp 243 - 244 - DOI: <https://doi.org/10.23637/ERADOC-1-157>

67/C/37.1

WINTER WHEAT

(EB)

Chemical control of take-all (*Ophiobolus graminis*) - Claycroft 1967.

Design: 3 randomised blocks of 20 plots.

Area of each plot: 0.0040. Area harvested: 0.0021.

Treatments: None, 125 lb kaolin dust applied (O) - two plots per block, and all combinations of:-

1. Chemicals:

DAC 2787 (tetrachloroisophthalonitrile)	(A)
Du Pont 1823 (1,4-dichloro-2,5-dimethoxybenzene)	(B)
Triphenyltin chloride	(C)
'Cela A36'	(D)
1-phenyl-3-thiosemicarbazide	(E)
Triamiphos, (5-amino-1-(bisdimethylaminophosphinyl)-3-phenyl-1,2,4 triazole)	(F)

2. Rates of application in lb:

A	B	C	D	E	F	
2.5	1.25	0.25	0.5	1.25	0.9	(R1)
10.0	5.0	1.0	2.0	5.0	1.8	(R2)
40.0	20.0	4.0	8.0	20.0	3.6	(R3)

Treatments A to E were made up with kaolin dust to the standard quantity for application at 125 lb. Treatment F was applied as a wettable powder spray at 70 gals.

Basal applications: 340 lb (6:15:15) combine drilled. 1.2 cwt N as 'Nitro-Chalk' in spring. Weedkiller: Aminotriazole at 4 lb and ammonium thiocyanate at 3.7 lb in 40 gals.

Cultivations, etc.: Weedkiller applied: Sept 20, 1966. Ploughed: Oct 6. Treatments A to E and O applied: Oct 17. Seed drilled at 165 lb: Oct 27. Treatment F applied: Mar 7, 1967. Basal 'Nitro-Chalk' applied: Apr 19. Combine harvested: Aug 30. Variety: Cappelle. Previous crops: Barley 1965 and 1966.

NOTE: Samples were taken in May and July for the estimation of take-all (*Ophiobolus graminis*), eyespot (*Cercospora herpotrichoides*), sharp eyespot (*Rhizoctonia solani*), and brown root rot.

Standard error per plot.

Grain: 3.29 or 7.9% (34 d.f.)

67/c/37.2

SUMMARY OF RESULTS

GRAIN

	A	B	C	D	E	F	Mean
	(±1.90)						(±0.77)
R1	42.1	43.5	39.8	41.5	42.5	42.5	42.0
R2	45.6	42.6	39.2	36.9	40.7	40.4	40.9
R3	41.6	40.1	42.1	41.5	45.0	42.8	42.2
Mean (±1.10)	43.1	42.1	40.4	40.0	42.7	41.9	41.7

O 43.1 (±1.34)

Mean D.M. % (all plots): 85.1