


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1967

[Full Table of Content](#)



67/W/WEC/C/42 Nematode Resistant Barley - Barley - Pathogen Infested Site

Rothamsted Research

Rothamsted Research (1968) *67/W/WEC/C/42 Nematode Resistant Barley - Barley - Pathogen Infested Site* ; Yields Of The Field Experiments 1967, pp 261 - 262 - **DOI:**
<https://doi.org/10.23637/ERADOC-1-157>

67/C/42.1

NEMATODE - RESISTANT BARLEY

(on site of 'Scorch' study)

(WEC)

Effect of using resistant and susceptible segregates of barley on land heavily infested with *Heterodera avenae* and the effect on yields - Woburn Butt Close 1967.

Design: Single replicate of 3 x 2 x 2 x 2 x 2 in blocks of 12 plots, with certain high order interactions confounded with block differences.

Area of each plot: 0.0032. Area harvested: 0.0021.

Treatments: All combinations of:-

1. Nitrogen: 0.4 (N1), 0.8 (N2), 1.2 cwt N (N3) as 'Nitro-Chalk' in the seedbed (N1, N2 and N3 being cumulative with N rates 1964 - 66)
2. Formalin 1964: None (O) Formalin (F64)
3. Formalin 1965: None (O) Formalin (F65)
4. Formalin 1966: None (O) Formalin (F66)
5. Varieties: Resistant segregate (R), susceptible segregate (S).

Basal applications: 20 cwt ground chalk 250 lb (0:20:20). Weedkillers: 4 lb amino-triazole plus 3.7 lb ammonium thiocyanate in 33 gals and ioxynil/mecoprop (Actril C at 5 pints in 50 gals).

Cultivations, etc.: Weedkiller applied: Sept 27, 1966. Ground chalk applied: Oct 31. Ploughed: Nov 2. PK compound applied by machine, 'Nitro-Chalk' applied by hand, seed drilled at 140 lb: Mar 15, 1967. Weedkiller (Actril C) applied: May 25. Combine harvested: Aug 23.

- NOTES (1) Crop samples for nematode counts and plant weights were taken in late May and early July.
- (2) Soil samples taken after harvest for nematode population counts.
- (3) For previous years' results see 'Results' 64/Da/3, 65/C/30 and 66/C/23.

Standard error per plot.

Grain: 5.19 or 24.5% (11 d.f.)

67/c/42.2

SUMMARY OF RESULTS

GRAIN

	R	S	Mean
Mean (± 1.06)	20.7 (± 1.84)	21.8	21.2 (± 1.30)
N1	14.7	14.6	14.7
N2	21.1	22.1	21.6
N3	26.2	28.6	27.4
	(± 1.50)		(± 1.06)
O	20.2	21.7	21.0
F64	21.1	21.8	21.5
	(± 1.50)		(± 1.06)
C	19.6	22.3	20.9
F65	21.7	21.3	21.5
	(± 1.50)		(± 1.06)
O	21.5	22.6	22.0
F66	19.8	20.9	20.4

Mean D.M. %: 80.0