


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1967

[Full Table of Content](#)



67/R/RW101/DA/1 Winter Wheat - Seed Rates, Methods of Sowing and Bulb Fly

Rothamsted Research

Rothamsted Research (1968) *67/R/RW101/DA/1 Winter Wheat - Seed Rates, Methods of Sowing and Bulb Fly* ; Yields Of The Field Experiments 1967, pp 275 - 276 - DOI:

<https://doi.org/10.23637/ERADOC-1-157>

67/Da/1.1

WINTER WHEAT

(RW 101)

Wheat bulb fly - seed rates and methods of sowing - Stackyard 1967.

Design: 6 randomised blocks of 4 plots.

Area of each plot: 0.0193. Area harvested: 0.0138.

Treatments: All combinations of:-

1. Sowing methods: Broadcast by drill (B), drilled (D).
2. Seed rates: 65 lb (R1), 183 (R3).

Basal applications: 280 lb (6:15:15) broadcast by drill. 0.84 cwt N as 'Nitro-Chalk' in spring. Weedkiller: Ioxynil/mecoprop (Atril C at 6 pints in 20 gals).

Cultivations, etc.: Rotary cultivated: Mar 30 and May 19, 1966.
Deep-tine cultivated: June 9. Rotary cultivated: July 1.
Seed sown, NPK basal compound applied: Nov 4. 'Nitro-Chalk' applied: Apr 18, 1967. Sprayed: May 9. Combine harvested: Aug 30. Variety: Cappelle. Previous crops: Winter wheat 1965, fallow 1966.

NOTE: Samples were taken from late February until mid-May to estimate numbers of plants, shoots, larvae and pupae and damaged plants and shoots.

Standard error per plot.

Grain: 3.45 or 10.2% (13 d.f.)

67/Da/1.2

SUMMARY OF RESULTS

GRAIN

	R1	R3	Mean
	(±1.41)		(±1.00)
B	29.5	41.0	35.3
D	20.8	43.5	32.1
Mean (±1.00)	25.2	42.2	33.7

Mean D.M. %: 84.7