

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1967

[Full Table of Content](#)



67/R/RW201/DA/2 Winter Wheat - Ccc and Eyespot

Rothamsted Research

Rothamsted Research (1968) *67/R/RW201/DA/2 Winter Wheat - Ccc and Eyespot* ; Yields Of The Field Experiments 1967, pp 277 - 278 - DOI: <https://doi.org/10.23637/ERADOC-1-157>

67/Da/2.1

WINTER WHEAT

(RW 201)

CCC and eyespot - Claycroft 1967.

Design: 4 x 4 Latin square.

Area of each plot: 0.0202. Area harvested: 0.0104.

Treatments: No chlormequat	(O)
Sprayed with chlormequat* at 2.5 lb in 37 gals:-	
Early (at 3 leaf stage)	(E)
At 5 leaf stage	(M)
Late (at shooting stage)	(L)

* 2-chloroethyltrimethylammonium chloride, formerly described as CCC.

Basal applications: 410 lb (6:15:15) combine drilled, 1.2 cwt N
as 'Nitro-Chalk' in spring.

Cultivations, etc.: Ploughed: Oct 6, 1966. Seed drilled at 180 lb:
Oct 28. Chlormequat applied: E - Feb 6, 1967 and M - Mar 21.
'Nitro-Chalk' applied: Apr 19. Chlormequat applied: L - Apr 26.
Combine harvested: Aug 31. Variety: Champlain. Previous crops:
Barley 1965 and 1966.

NOTE: Samples were taken on 29th June for estimation of eyespot
(*Cercospora herpotrichoides*).

Standard error per plot.
Grain: 2.45 or 4.2% (15 d.f.)

67/Da/2.2

SUMMARY OF RESULTS

GRAIN

O	E	M	L	Mean
54.9	60.4	58.0	58.6	58.0
	(±1.23)			

Mean D.M. %: 83.2