


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1967

[Full Table of Content](#)



67/W/WW101/DA/9 Spring Wheat - Ccc and Irrigation

Rothamsted Research

Rothamsted Research (1968) *67/W/WW101/DA/9 Spring Wheat - Ccc and Irrigation* ; Yields Of The Field Experiments 1967, pp 293 - 295 - DOI: <https://doi.org/10.23637/ERADOC-1-157>

67/Da/9.1

SPRING WHEAT

(WW 101)

CCC, irrigation, and nitrogen - Woburn Butt Close (Series II) 1967.

Design: 4 blocks of 2 plots split into 2 for CCC and again into 2 for nitrogen N₄ - N₃ + N₂ - N₁ on half plots.

Area of each quarter plot: 0.0136. Area harvested: 0.0036.

Treatments: All combinations of:-

Main plots: 1. Irrigation: None (O), full irrigation (C).

Half plots: 2. Chlormequat*: None (O), chlormequat sprayed at 2.5 lb in 37 gals (S).

Quarter plots: 3. Nitrogen: 0.4 (N₁), 0.8 (N₂), 1.2 (N₃), 1.6 (N₄) cwt N as 'Nitro-Chalk'.

* 2-chloroethytrimethylammonium chloride, formerly described as CCC.

Basal applications: 240 lb (0:14:28) combine drilled. Weedkiller: Ioxynil/mecoprop (Actril C at 5 pints in 25 gals).

Cultivations, etc.: Ploughed: Nov 3, 1966. Basal PK applied, seed drilled at 170 lb: Mar 7, 1967. 'Nitro-Chalk' applied: Mar 14. Weedkiller applied: Apr 28. CCC applied: May 23. C plots, irrigation applied at 0.25 inches: July 11, July 21, at 0.5 on each of the following occasions: June 12, June 15, July 4, July 10, July 17, and at 1.0 inch: June 20 (total 4 inches). Combine harvested: Aug 24. Variety: Klokka.

- NOTES: (1) Plant samples were taken on July 5 for estimates of dry weight of tops and roots. Soil cores were also taken for estimates of the weight of roots left in the soil.
(2) Sub-plots 101 - 132 were sampled on Aug 22 for estimates of straw height and weight.

Standard errors per plot. Grain:

1/2 plot: 2.38 or 5.3% (4 d.f.)

1/4 plot: 2.56 or 5.7% (8 d.f.)

67/Da/9.2

SUMMARY OF RESULTS

GRAIN

	N1	N2	N3	N4	Mean
	(1) and (2)				
Mean	37.5	49.4	48.4	44.1	44.8
	(3) and (4)				
O	35.7	47.2	42.9	38.4	41.0
C	39.2	51.5	53.9	49.9	48.6
	(5) and (6)				(± 0.84)
O	37.6	48.7	49.7	42.9	44.7
S	37.3	50.0	47.1	45.4	45.0

Mean D.M. %: 80.6

- (1) (± 0.90) For use in comparisons involving N4 - N2 or N3 - N1
 (2) (± 1.06) For use in all other comparisons
 (3) (± 1.28) For use in horizontal and interaction comparisons
 (4) (± 1.68) For use in all other horizontal and interaction comparisons
 (5) (± 1.28) For use in horizontal and interaction comparisons involving
 N2 - N2 or N3 - N1
 (6) (± 1.60) For use in all other comparisons

67/Da/9.3

	STRAW				
	N1	N2	N3	N4	Mean
Mean	33.4	50.3	54.9	57.8	49.1
O	31.6	47.8	50.0	55.8	46.3
C	35.2	52.8	59.8	59.8	51.9
O	37.6	53.1	61.6	62.7	53.7
S	29.2	47.5	48.3	52.9	44.5

Mean D.M. %: 83.2