


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1967

[Full Table of Content](#)



67/R/RP8/1/DD/7 Potatoes - Oospora Free Seed

Rothamsted Research

Rothamsted Research (1968) *67/R/RP8/1/DD/7 Potatoes - Oospora Free Seed* ; Yields Of The Field Experiments 1967, pp 327 - 328 - DOI: <https://doi.org/10.23637/ERADOC-1-157>

67/Dd/7.1

POTATOES

(RP 8/1)

Oospora*-free seed - Pastures 1967.

Design: 6 blocks of 3 plots.

Area of each plot: 0.0071. Area harvested: 0.0031.

Treatments: Oospora-free seed (F), Oospora-free seed re-infected with Oospora (R), normal stock seed (N).

* Oospora pustulans - skinspot.

Basal applications: 10 cwt (13:13:20). Weedkiller: Linuron at 0.75 lb plus paraquat at 0.5 lb ion in 37 gals. Fungicide: Mancozeb at 1.2 lb in 30 gals on four occasions.

Cultivations, etc.: - Rotary cultivated: Sept 16, 1966. Ploughed: Nov 23 - Jan 5, 1967. Basal NPK compound applied: Mar 30. Rotary cultivated second time, potatoes planted: Apr 24. Weedkiller applied: May 18. Fungicide applied: June 30, July 21, Aug 3, Aug 24. Sprayed with undiluted BOV at 15 gals: Sept 20. Haulm destroyed mechanically, potatoes lifted: Sept 28. Variety: King Edward. Previous crops: Spring wheat and barley 1965, fallow 1966.

Standard error per plot:

Total tubers: 1.642 or 6.9% (10 d.f.)

67/Da/7.2

SUMMARY OF RESULTS

F	R	N	Mean
TOTAL TUBERS			
	(± 0.670)		
24.74	22.80	23.78	23.77
% WARE			
78.7	87.6	86.4	84.2