

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1967

[Full Table of Content](#)



67/R/CW/B/7 Cultivation-weedkiller Rotation

Rothamsted Research

Rothamsted Research (1968) *67/R/CW/B/7 Cultivation-weedkiller Rotation ; Yields Of The Field Experiments 1967*, pp 99 - 103 - DOI: <https://doi.org/10.23637/ERADOC-1-157>

67/B/7.1

CULTIVATION - WEEDKILLER ROTATION

(CW)

Great Harpenden 1967 - the 7th year

A comparison of weed-control by various cultivation methods and by pre-emergence weedkillers.

For previous history, rotations, treatments etc., see 'Results' 61/B/10, 62/B/10, 63/B/10, 64/B/9, 65/B/8, 66/B/7.

Area harvested: Beans - 0.0100, wheat, potatoes and barley - 0.0107.

Revised basal dressings:-

Potatoes: 10 cwt (13:13:20).

Spring wheat: 3 cwt (25:10:10).

Barley: 3 cwt (25:10:10).

Wheat and barley: Basal post-emergence weedkiller was applied in 1967, the test of none v. weedkiller being omitted because of unusually large quantities of weeds.

NOTE: After the potato crop of 1966 all plots were spring-time cultivated twice in autumn and once in spring. The resulting tilth on the A and B plots (which had had no other cultivation) was considered adequate and no other cultivations were done before sowing barley.

Cultivations, etc.:-

Spring beans: T plots deep-tine cultivated, depth 3-4 ins:

Nov 5, 1966. T plots deep-tine cultivated, depth 5-6 ins:

Nov 11. P and C plots ploughed: Nov 15. R plots rotary cultivated, depth 6 ins: Nov 17. B plots sprayed with paraquat at 0.75 lb ion in 40 gals: Nov 21. B plots deep-tine cultivated, depth 4-5 ins: Nov 29. P, C and T plots spring-time cultivated: Mar 4, 1967. P, C, T and B plots spring-time cultivated:

Mar 7. A plots rotary cultivated, seed drilled at 200 lb, S plots sprayed: Mar 8. M and C plots tractor-hoed twice: May 2 and June 2. Sprayed with demeton-s-methyl (Metasystox at 12 fluid oz in 37 gals): June 13 and July 7 (the second spray to edges of blocks only). Sprayed by aircraft with demeton-s-methyl (Metasystox at 12 fluid oz in 3 gals): July 11. Combine harvested: Sept 1. Variety: Tarvin.

Spring wheat: Sprayed with aminotriazole at 4 lb and ammonium thiocyanate at 3.7 lb in 40 gals: Sept 19, 1966. T plots deep-tine cultivated, depth 3-4 ins: Nov 5, 1966. T plots deep-tine cultivated, depth 5-6 ins: Nov 12. P and C plots ploughed: Nov 14. R plots rotary cultivated, depth 6 ins: Nov 17. Plots spring-time cultivated - B plots: Nov 29, P, T, C and B plots:

Mar 3, 1967, P and C plots: Mar 7, P, T, R and C plots: Mar 15. A plots rotary cultivated, seed drilled at 175 lb:

67/B/7.2

Mar 15. All plots rolled: Apr 18. All plots sprayed with mecoprop/2,4-D (Methoxone Extra at 6 pints in 30 gals):
 May 12. Combine harvested: Aug 24. Variety: Kloka.
 Potatoes: T plots deep-tine cultivated, depth 3-4 ins: Nov 5, 1966.
 T plots deep-tine cultivated, depth 5-6 ins: Nov 12. P and C plots ploughed: Nov 14. R plots rotary cultivated: Nov 18.
 P, T and C plots deep-tine cultivated: Mar 20, 1967. Basal compound fertiliser applied: Mar 22. P, T and C plots spring-tine cultivated twice: Mar 29 and Apr 3. R, A and B plots rotary cultivated, potatoes machine planted: Apr 4. Ridges rolled: Apr 28. S plots sprayed: May 9. M and C plots grubbed: May 12. M plots mechanically weeded twice: May 13.
 C plots mechanically weeded twice: May 19. M and C plots grubbed: June 14. All plots sprayed with mancozeb at 1.2 lb in 30 gals: June 29. M, C and Y plots rotary ridged: June 30. All plots sprayed three times with mancozeb at 1.2 lb in 30 gals: July 20, Aug 4, Aug 24. Sprayed with undiluted BOV at 15 gals: Sept 6. Haulm destroyed mechanically: Sept 13. Lifted: Sept 14. Variety: Pentland Dell.
 Barley: All plots sprayed with sodium trichloroacetate at 18 lb in 40 gals: Oct 22, 1966. All plots spring-tine cultivated: Nov 1. All plots sprayed with sodium trichloroacetate at 18 lb in 40 gals: Nov 21. All plots spring-tine cultivated: Nov 29. T plots deep-tine cultivated, first stroke at 20, second at 40, depth 6-7 ins, P and C plots ploughed: Jan 17, 1967. R plots rotary cultivated, depth 6 ins: Feb 6. All plots spring-tine cultivated: Mar 4. P, T and C plots spring-tine cultivated, seed drilled at 140 lb: Mar 6. All plots rolled: Apr 18. All plots sprayed with mecoprop/2,4-D (Methoxone Extra at 6 pints in 30 gals): May 12. Combine harvested: Aug 21. Variety: Maris Badger.

Standard errors per whole plot.

Spring beans.	Grain:	2.93 or 9.6% (8 d.f.)
Wheat.	Grain:	6.22 or 15.7% (8 d.f.)
Potatoes.	Total tubers:	1.887 or 13.7% (8 d.f.)
Barley.	Grain:	1.94 or 4.2% (8 d.f.)

67/B/7.3

SUMMARY OF RESULTS

	P	R	T	Mean
--	---	---	---	------

SPRING BEANS

GRAIN

Mean (± 1.20)	32.5	29.7	29.9	30.7
M (± 2.08)	33.1	28.8	29.8	30.6 (± 1.20)
S (± 1.47)	32.2	30.1	30.0	30.8 (± 0.85)

A

B

C

32.9

31.9

30.6

General mean: 31.0

Mean D.M. %: 81.6

SPRING WHEAT

GRAIN

Mean (± 2.54)	37.5	41.6	40.1	39.7
M (± 4.40)	34.6	39.2	41.0	38.3 (± 2.54)
S (± 3.11)	38.9	42.8	39.6	40.5 (± 1.80)

A

B

C

42.4

45.8

43.6

General mean: 40.8

Mean D.M. %: 82.2

				67/B/7.4
	P	R	T	Mean
POTATOES				
TOTAL TUBERS				
Mean (± 0.770)	14.76	13.35	13.11	13.74
		(± 1.334)		(± 0.770)
M	14.38	12.09	11.20	12.56
S	13.89	14.23	14.45	14.19
SY	16.00	13.73	13.68	14.47
	A	B	C	
	15.50	14.50	12.23	
General mean: 13.82				
% WARE				
Mean	97.7	96.5	96.2	96.8
M	97.9	97.2	96.8	97.3
S	97.4	96.5	94.2	96.1
SY	97.9	95.8	97.6	97.1
	A	B	C	
	97.4	96.1	97.5	
General mean: 96.8				

67/B/7.5

BARLEY

GRAIN

	P	R	T	Mean
Mean (± 0.79)	46.6	45.6	47.1	46.4
M	47.0	(± 1.38) 43.8	46.8	(± 0.79) 45.9
S	46.1	46.6	48.3	47.0
SY	46.7	46.4	46.1	46.4

A

B

C

44.4

45.6

45.0

General mean: 46.1

Mean D.M. %: 80.4