

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1968

[Full Table of Content](#)



68/C/2 Intensive Barley Growing Experiment - S. Wheat, Barley, Fallow

Rothamsted Research

Rothamsted Research (1970) *68/C/2 Intensive Barley Growing Experiment - S. Wheat, Barley, Fallow* ; Numerical Results Of The Field Experiments 1968, pp 137 - 138 - DOI:
<https://doi.org/10.23637/ERADOC-1-58>

68/c/2.1

INTENSIVE BARLEY GROWING EXPERIMENT

(IB)

Little Knott I - 1968, the eighth year

For treatments, etc., see 'Results' 61/C/8 (N0 = None, N1 = 0.3, N2 = 0.6, N3 = 0.9 cwt N).

Area of each plot: 0.0212. Area harvested: Spring wheat - 0.0188, barley - 0.0139.

Winter wheat, sequence 9: These plots were fallowed in 1968 because of Blackgrass (*Alopecurus myosuroides*) and Couchgrass (*Agropyron repens*).

Basal applications: Manures as previously.

Weedkillers: All plots: Aminotriazole at 4 lb and ammonium thiocyanate at 3.7 lb in 32 gals. Spring wheat and barley: Ioxynil at 7.5 oz and mecoprop at 22.5 oz in 20 gals.

Cultivations, etc.: Aminotriazole etc. applied: 21 Sept, 1967.

Ploughed: 30 Oct.

Spring wheat: Seed combine drilled at 175 lb: 6 Mar, 1968.

'Nitro-Chalk' applied: 20 Mar. Weedkiller applied: 6 May.

Combine harvested: 24 Aug.

Barley: Seed combine drilled at 140 lb: 6 Mar, 1968. 'Nitro-Chalk' applied: 20 Mar. Weedkiller applied: 6 May. Combine harvested: 20 Aug.

Fallow plots: Rotary cultivated 4 times: 12 Apr, 29 May, 18 June, 6 Aug, 1968.

- NOTES: (1) Yields were taken for all sequences except 9 (fallow).
(2) Estimates of eyespot (*Cercospora herpotrichoides*) and take-all (*Ophiobolus graminis*) were made in spring and summer.
(3) For previous years' results see 'Results' 61/C/8, 62/C/7, 63/C/2, 64/C/2, 65/C/2, 66/C/2, 67/C/2.

Standard errors per plot.

Barley, Grain: 2.86 or 9.6% (31 d.f.)

68/c/2.2

SUMMARY OF RESULTS

SPRING WHEAT (8)

CROP 1961 - 1967 SW

GRAIN				
NO	N1	N2	N3	Mean
11.4	13.8	20.3	19.9	16.3

Mean D.M. %: 82.1

BARLEY (1 - 7 & 10)

CROP IN								GRAIN				
61	62	63	64	65	66	67		NO	N1	N2	N3	Mean
								(±2.02)				(±1.01)
O	Be	B	B	B	B	B		12.8	27.1	33.1	35.2	27.1
SW	O	Be	B	B	B	B		16.4	24.2	32.5	35.1	27.1
O	SW	O	Be	B	B	B		15.3	25.1	30.5	34.3	26.3
Be	O	SW	O	Be	B	B		14.0	27.3	31.5	35.4	27.0
SW	Be	O	SW	O	Be	B		26.9	29.7	34.0	36.2	31.7
SW	SW	Be	O	SW	O	Be		30.3	36.3	38.6	35.9	35.3
B	B	B	B	B	B	B		15.0	28.0	35.9	36.6	28.9
Be	WW	P	B	Be	WW	P		32.2	35.9	35.5	34.4	34.5
Mean (±0.71)								20.4	29.2	33.9	35.4	29.7

Mean D.M. %: 82.7