

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1968

[Full Table of Content](#)



68/C/4 Residues of N, P and K to Grass - Spring Wheat

Rothamsted Research

Rothamsted Research (1970) *68/C/4 Residues of N, P and K to Grass - Spring Wheat* ; Numerical Results Of The Field Experiments 1968, pp 145 - 146 - **DOI:**

<https://doi.org/10.23637/ERADOC-1-58>

68/C/4.1

SPRING WHEAT

(AF)

Residues of N,P and K to grass (the 11th year of 'Levels of N and K to grass'), Harwoods Piece 1968.

For treatments etc. see 'Results' 63/C/7.1 and 65/C/6.2 and for previous years' results see 58/Cg/2, 59/Cg/2, 60/Ci/1, 61/Dg/1, 62/C/11, 63/C/7, 64/C/6, 65/C/6, 66/C/5 and 67/C/4.

Area of each plot: 0.0082. Area harvested: 0.0054.

No treatment or basal P and K were applied in 1968.

Basal applications: 1.0 cwt N as 'Nitro-Chalk' broadcast by drill.
Weedkiller: Ioxynil at 7.5 oz and mecoprop at 22.5 oz in 20 gals.

Cultivations, etc.: Ploughed: 8 Nov, 1967. Seed drilled at 185 lb, 'Nitro-Chalk' applied: 5 Mar, 1968. Weedkiller applied: 15 May. Combine harvested: 27 Aug. Variety: Klokka.

NOTE: Soil samples were taken in October 1967 for pH and available P and K. Crop samples were taken at ear emergence for yield of dry matter and for determination of percentages of N, P and K. Grain and straw samples were taken at harvest for determination of percentages of N, P and K.

Standard error per plot:
Grain: 2.86 or 11.8% (33 d.f.)

68/c/4.2

SUMMARY OF RESULTS

N	0	1	1	1	2	2	2	3	3	3	3	3	Mean
P	1	1	1	1	1	1	1	1	1	1	0	2	
K	0	0	1	2	0	1	2	0	1	2	2	2	

GRAIN

(± 1.43)

31.0 20.6 29.8 31.3 15.8 22.9 27.0 16.5 21.9 23.5 25.4 24.2 | 24.2

STRAW

33.3 20.9 33.5 39.9 20.5 27.1 33.6 14.9 21.6 28.8 32.5 32.1 | 28.2

Mean D.M. %: Grain: 82.5
Straw: 59.8

NOTE: Treatments were applied 1958/67