

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1968


[Full Table of Content](#)



68/C/24 the Effect of Granule Size on the Rate of Release of N from Ibdur - Grass

Rothamsted Research

Rothamsted Research (1970) *68/C/24 the Effect of Granule Size on the Rate of Release of N from Ibdur - Grass* ; Numerical Results Of The Field Experiments 1968, pp 207 - 210 - **DOI:**
<https://doi.org/10.23637/ERADOC-1-58>



68/c/18.3

LEY: DRY MATTER

LC	LN
1ST CUT	
23.1	26.8
2ND CUT	
16.5	25.2
TOTAL OF 2 CUTS	
39.6	52.0

Mean D.M. %: 1st cut: 21.7
2nd cut: 21.4
Total of 2 cuts: 21.4

68/c/18.2

SUMMARY OF RESULTS

WINTER WHEAT

	NO	N1	N2	N3	Mean
	GRAIN				
	(1) and (2)				(±1.09)
DG	30.3	39.8	39.0	35.6	36.2
ST	20.5	31.2	36.4	38.2	31.6
PT	21.1	30.4	36.5	37.5	31.4
GM	25.5	37.7	36.0	35.5	33.7
FD	20.6	33.4	37.5	36.8	32.1
FS	21.0	31.6	37.0	38.4	32.0
Mean (±0.57)	23.2	34.0	37.1	37.0	32.8

(1) (±1.63) For use in vertical and diagonal comparisons only

(2) (±1.39) For use in horizontal and interaction comparisons only

STRAW

DG	32.9	44.3	46.8	49.9	43.5
ST	21.9	35.2	42.3	46.0	36.3
PT	24.0	35.0	42.6	43.5	36.3
GM	20.3	35.1	41.3	41.7	34.6
FD	22.0	37.7	41.0	42.2	35.7
FS	24.7	36.1	42.3	43.1	36.5
Mean	24.3	37.2	42.7	44.4	37.2

Mean D.M. %: Grain: 81.1
Straw: 79.2

68/c/18.1

ORGANIC MANURING EXPERIMENT

(WOM)

The cumulative effects of organic matter on light land - Woburn Stackyard B 1968, 4th year.

For previous history, rotation, treatments etc., see 'Results' 66/c/31 and 67/c/24. All plots except those under leys (LC and LN) carried winter wheat.

Area of each sub plot: 0.0156. Area harvested: Wheat - 0.0104
leys - 0.0129.

Fertilisers applied Autumn 1967 (cwt)

Treatment	P2O5	K2O	MgO
DG	-	-	-
ST	0.4	-	0.15
PT	0.5	1.0	-
GM	0.5	1.4*	0.2
FD	2.2	1.6	0.8
FS	0.5	1.0	0.2
LC	0.5	2.0	0.2
LN	0.5	2.0	0.2

* Including 0.4 applied in spring before undersowing.

Nitrogen to winter wheat:

NO, N1, N2, N3. 0.0, 0.4, 0.8, 1.2 as 'Nitro-Chalk' in addition to basal application of 0.2 cwt N.

Cultivations, etc.:

LC and LN plots: P, K, and Mg applied: 3 Oct, 1967.

'Nitro-Chalk' applied to LN plots: 27 Mar, 1968, 14 June.

Cut twice: 5 June, 10 Oct.

Wheat: Subsoiled: 25 Sept, 1967. PK and Mg applied: 10 Oct.

Peat, straw and FYM applied, ploughed: 11 Oct. Seed

drilled at 190 lb: 25 Oct. K applied to GM plots: 27 Mar, 1968.

GM plots undersown with late flowering red clover at 15 lb:

29 Mar. N applied: 18 Apr. All plots except GM plots

sprayed with ioxynil/mecoprop (Actril C at 6 pints in

25 gals): 25 Apr. Combine harvested: 23 Aug. Variety: Cappelle.

Standard errors per plot. Winter wheat:

Whole plot: 2.19 or 6.7% (15 d.f.)

Sub plot: 2.79 or 8.5% (54 d.f.)