

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1968

[Full Table of Content](#)



68/A/4 Agdell Grass

Rothamsted Research

Rothamsted Research (1970) *68/A/4 Agdell Grass* ; Numerical Results Of The Field Experiments 1968, pp 31 - 34 - DOI: <https://doi.org/10.23637/ERADOC-1-58>

68/A/4.1

GRASS - AGDELL 1968

(AG)

For history, treatments, etc. see 'Details' 1962 and 'Results' 63/A/4, 64/A/4, 65/A/4, 66/A/4 and 67/A/4.

Area of each microplot: Plots 1 - 4 - 0.0180. Plots 5 - 6 - 0.0162.
Area harvested: 1st cut: Plots 1 - 4 - 0.0046. Plots 5 - 6 - 0.0040. 2nd cut: Plots 1 - 4 - 0.0023. Plots 5 - 6 - 0.0020.

P (as triple superphosphate) and K (as muriate of potash) were applied in March 1968 to balance removals by grass in 1967 to all sub plots except P0, which continues to receive no P, and K0, which continues to receive no K.

Rates in cwt P2O5

Plot no.	Sub plots testing P:-				Sub plots testing K:-			
	P0	P1	P2	P4	K0	K1	K2	K4
1	0	0.36	0.32	0.35	0.23	0.36	0.40	0.35
2	0	0.36	0.40	0.31	0.22	0.38	0.39	0.40
3	0	0.39	0.32	0.42	0.18	0.32	0.31	0.32
4	0	0.28	0.27	0.33	0.31	0.36	0.34	0.31
5	0	0.30	0.31	0.30	0.33	0.38	0.38	0.35
6	0	0.30	0.31	0.33	0.22	0.33	0.34	0.33

Rates in cwt K2O

Plot no.	Sub plots testing P:-				Sub plots testing K:-			
	P0	P1	P2	P4	K0	K1	K2	K4
1	1.33	1.80	1.56	1.65	0	1.48	1.80	1.66
2	0.86	1.63	1.99	1.48	0	1.62	1.68	1.84
3	0.99	2.00	1.57	1.99	0	1.38	1.35	1.52
4	0.48	1.68	1.67	1.75	0	1.50	1.67	1.54
5	0.38	1.84	2.04	1.86	0	1.96	1.94	2.00
6	0.18	1.68	1.57	1.72	0	1.53	1.54	1.64

Basal dressing: 'Nitro-Chalk' applied at 0.8 cwt N on 8 Sept, 1967 and 15 Mar, 1968, 30 May, 15 July.

Cultivations, etc.:

Grass: Ground chalk applied at 46 cwt to plots 1 and 2 and to the southern half of plots 3 and 4: 6 June, 1967. Ploughed: 14 June. Rotary cultivated 4 times: 5 July, 21 July, 8 Aug, 24 Aug. Seed

pp 3

68/A/4.3

SUMMARY OF RESULTS

DRY MATTER

Plot

P K	5	6	3	4	1	2	Mean
1ST CUT							
0 4	0.0	0.0	3.0	0.0	17.9	5.9	4.5
1 4	20.8	27.6	37.9	31.3	38.1	34.6	31.7
2 4	28.8	27.5	33.0	30.7	33.6	38.6	32.0
4 4	29.8	29.4	36.5	30.7	35.8	40.5	33.8
4 0	17.1	26.0	29.1	29.3	26.5	22.8	25.1
4 1	26.7	31.5	40.3	33.5	35.6	35.7	33.9
4 2	30.5	34.8	31.2	35.6	39.4	35.0	34.4
4 4	31.1	30.8	37.1	39.4	37.7	38.5	35.8
Mean	23.1	25.9	31.0	28.8	33.1	31.5	28.9

Mean D.M. %: 15.0

2ND CUT							
0 4	3.7	4.8	15.5	8.0	23.7	18.7	12.4
1 4	17.7	16.8	16.2	17.0	22.6	20.3	18.4
2 4	17.9	16.9	19.5	13.1	21.1	22.1	18.4
4 4	17.4	16.8	16.0	20.1	22.0	19.7	18.7
4 0	21.1	15.8	17.8	21.4	17.1	17.1	18.4
4 1	19.0	17.1	13.5	25.8	21.2	23.9	20.1
4 2	20.6	18.2	14.7	15.2	18.6	22.8	18.4
4 4	23.2	15.0	16.0	17.3	20.3	20.1	18.6
Mean	17.6	15.2	16.2	17.2	20.8	20.6	17.9

Mean D.M. %: 15.6

68/A/4.4

DRY MATTER

P K	Plot						Mean
	5	6	3	4	1	2	
	3RD CUT						
0 4	19.6	21.3	26.2	24.8	23.1	21.4	22.7
1 4	30.7	28.5	24.4	35.2	26.1	27.8	28.8
2 4	25.3	37.0	28.2	31.8	20.5	27.8	28.4
4 4	25.8	27.0	25.6	24.6	26.2	25.4	25.8
4 0	18.2	23.4	20.2	21.0	14.2	13.1	18.3
4 1	25.8	26.1	22.4	28.2	23.7	24.6	25.1
4 2	24.5	29.5	25.4	27.8	27.9	28.8	27.3
4 4	26.1	26.7	23.2	29.3	24.3	25.7	25.9
Mean	24.5	27.4	24.4	27.8	23.2	24.3	25.3

Mean D.M. %: 20.7

TOTAL OF 3 CUTS							
0 4	23.3	26.2	44.6	32.8	64.7	46.0	39.6
1 4	69.3	72.9	78.5	83.4	86.8	82.7	78.9
2 4	72.0	81.4	80.7	75.6	75.2	88.4	78.9
4 4	73.0	73.2	78.1	75.3	84.0	85.6	78.2
4 0	56.4	65.3	67.1	71.6	57.8	53.0	61.9
4 1	71.5	74.6	76.2	87.5	80.5	84.2	79.1
4 2	75.7	82.5	71.3	78.6	85.9	86.7	80.1
4 4	80.3	72.4	76.3	86.1	82.4	84.3	80.3
Mean	65.2	68.6	71.6	73.9	77.1	76.4	72.1

Mean D.M. %: 17.1