

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1968

[Full Table of Content](#)



68/DB/1 Effects of Paths and Blank Rows - Barley

Rothamsted Research

Rothamsted Research (1970) *68/DB/1 Effects of Paths and Blank Rows - Barley* ; Numerical Results Of The Field Experiments 1968, pp 313 - 314 - DOI: <https://doi.org/10.23637/ERADOC-1-58>

68/Db/1.1

BARLEY

(RB 101)

Effects of paths and blank rows - West Barnfield I 1968.

Design: 4 randomised blocks of 8 plots.

Area of each plot: 0.0161. Area harvested: 0.0107.

Treatments: All combinations of:-

- | | |
|---|------|
| 1. Blank rows: None, 16 middle rows harvested | (R0) |
| 2 blank rows, 3(1) 16(1) 3 sown | (R1) |
| 4 blank rows, 2(2) 16(2) 2 sown | (R2) |
| 6 blank rows, 1(3) 16(3) 1 sown | (R3) |
| 2. N: 0.5 (N1), 1.0 (N2) cwt side placed as 'Nitro-Chalk. | |

Plots were of 24 rows each, 16 of which were harvested. (1) etc.
indicate number and position of blank (unsown) rows.

Basal applications: 280 lb (0:20:20) broadcast in the seedbed.

Weedkiller: Paraquat at 0.75 lb ion in 32 gals to stubble of
preceding crop.

Cultivations, etc.: Weedkiller applied: 6 Oct, 1967. Ploughed: 23 Oct.
Seed drilled at 145 lb, 'Nitro-Chalk' sideplaced, basal PK
applied: 29 Feb, 1968. Combine harvested: 21 Aug. Variety:
Maris Badger. Previous crops: Grass 1966, winter wheat 1967.

Standard error per plot.

Grain: 1.41 or 4.7% (21 d.f.)

68/Db/1.2

SUMMARY OF RESULTS

GRAIN (101 58)

	R0	R1	R2	R3	Mean
	(±0.71)				(±0.35)
N1	29.3	32.2	35.6	33.2	32.6
N2	26.8	27.2	28.8	29.4	28.0
Mean (±0.50)	28.1	29.7	32.2	31.3	30.3

Mean D.M. %: 81.7