

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1968

[Full Table of Content](#)



68/DB/5 Woburn Varieties and N - Barley

Rothamsted Research

Rothamsted Research (1970) *68/DB/5 Woburn Varieties and N - Barley* ; Numerical Results Of The Field Experiments 1968, pp 321 - 322 - DOI: <https://doi.org/10.23637/ERADOC-1-58>

68/Db/5.1

BARLEY

(WB 101)

Varieties and N - Woburn Great Hill Bottom I, 1968.

Design: 4 blocks of 4 plots, split for N.

Area of each sub plot: 0.0024. Area harvested: 0.0013.

Treatments: All combinations of:-

Whole plots: 1. Varieties: Impala (I), Maris Badger (MB),
Sultan (S), Zephyr (Z).

Sub plots: 2. Nitrogen: 0.6 (N2), 0.9 (N3), 1.2 (N4) cwt
N as 'Nitro-Chalk'.

Basal applications: 4 cwt (15:15:15) combine drilled. Weedkillers:
Paraquat at 0.5 lb ion in 33 gals, ioxynil/mecoprop (Actril
C at 5 pints in 25 gals).

Cultivations, etc.: Deep-tine cultivated: 25 Sept, 1967. Ploughed:
20 Feb, 1968. Paraquat applied: 2 Mar. Seed combine drilled at
140 lb: 7 Mar. 'Nitro-Chalk' applied: 14 Mar. Ioxynil/mecoprop
applied: 3 May. Combine harvested: 22 Aug. Previous crops:
Fallow 1966, potatoes 1967.

Standard errors per plot.

Grain: Whole plot: 3.14 or 11.9% (9 d.f.)

Sub plot: 4.10 or 15.6% (24 d.f.)

68/Db/5.2

SUMMARY OF RESULTS

	GRAIN				Mean
	I	MB	S	Z	
	(1) and (2)				(±1.03)
N2	25.4	22.7	30.2	28.3	26.6
N3	23.7	23.9	40.8	24.2	28.1
N4	20.7	20.6	32.0	22.8	24.0
Mean (±1.57)	23.3	22.4	34.3	25.1	26.3

- (1) (±2.29) For use in horizontal and diagonal comparisons only
 (2) (±2.05) For use in vertical and interaction comparisons only

Mean D.M. %: 82.9