


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1969

[Full Table of Content](#)



69/R/CS/24 - P, K & Take-all - Barley

Rothamsted Research

Rothamsted Research (1970) *69/R/CS/24 - P, K & Take-all - Barley* ; Numerical Results Of The Field Experiments 1969, pp 186 - 187 - DOI: <https://doi.org/10.23637/ERADOC-1-96>

BARLEY

(69/R/CS/24)

P, K and take-all (*Ophiobolus graminis*), West Barnfield II 1969, the second year. For treatments etc. and for the previous year's results see 'Results' 68/C/16.

Area of each plot: 0.0265. Area harvested: 0.0140.

Basal applications: 0.8 cwt N as 'Nitro-Chalk' combine drilled. Ground chalk at 23 cwt. Weedkillers: Paraquat at 0.75 lb ion in 20 gals, 2,4-D at 8 oz and dichlorprop at 32 oz in 20 gals.

Cultivations, etc.: Paraquat applied: 1 Oct, 1968. Ground chalk applied: 1 Nov. Ploughed: 19 Nov. P and K applied: 11 Mar, 1969. Seed combine drilled at 140 lb: 24 Mar. 2,4-D/dichlorprop applied: 20 May. Combine harvested: 22 Aug.

NOTE: Estimates of eyespot (*Cercospora herpotrichoides*) and take-all (*Ophiobolus graminis*) were made in spring and summer.

Standard error per plot.

Grain, cwt: 1.61 or 3.9% (27 d.f.)

SUMMARY OF RESULTS

	P0	P1	P4	P6*	P24*	Mean
	GRAIN:CWT					
	(±0.81)					(±0.36)
K1	33.1	42.0	43.4	43.0	42.8	40.9
K4	35.8	42.0	42.4	43.2	42.3	41.1
Mean (±0.57)	34.5	42.0	42.9	43.1	42.5	41.0

Mean D.M.%: 82.3

	STRAW:CWT					
K1	25.2	32.6	40.7	35.0	42.2	35.1
K4	25.3	35.8	42.9	38.2	42.2	36.9
Mean	25.3	34.2	41.8	36.6	42.2	36.0

Mean D.M.%: 92.0

* Residual