


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1969

[Full Table of Content](#)



69/W/CS/28 - Fumigants & Irrigation - Barley

Rothamsted Research

Rothamsted Research (1970) *69/W/CS/28 - Fumigants & Irrigation - Barley* ; Numerical Results Of The Field Experiments 1969, pp 201 - 202 - DOI: <https://doi.org/10.23637/ERADOC-1-96>

FUMIGANTS AND IRRIGATION

(69/W/CS/28)

The residual effects of fumigants for the control of *Pratylenchus* spp. in barley - Woburn Butt Close Series III, 2nd year. For design, treatments etc. and previous year's results see 'Results' 68/C/34.

Area of each sub plot: 0.0545. Area harvested: 0.0020.

Treatments: All combinations of:-

- Whole plots: 1. Irrigation: None (O), full irrigation (I).
- Quarter plots: 2. Residues of fumigants 1968: None (O), chloropicrin (C), D-D (D), Temik (T).
- Eighth plots: 3. Nitrogen in 1967 and 1968: None (NO), 0.4 (N1), 0.8 (N2), 1.2 (N3) cwt N as 'Nitro-Chalk'.

Basal applications: 280 lb (0:20:20). Weedkiller: Ioxynil at 7.5 oz and mecoprop at 22.5 oz in 25 gals.

Cultivations, etc.: Ploughed: 12 Nov, 1968. Seed combine drilled at 140 lb: 11 Mar, 1969. 'Nitro-Chalk' applied: 4 Apr. Weedkiller applied: 14 May. Irrigation applied at 0.5 inches: June 13, and 0.5 inches: July 4. Combine harvested: 11 Aug. Variety: Maris Badger.

NOTE: Crop and soil samples were taken at intervals through the growing season for nematode counts.

Standard errors per plot. Grain, cwt.

- Whole plot: 1.62 or 5.2% (6 d.f.)
- Quarter plot: 2.21 or 7.0% (24 d.f.)
- Eighth plot: 3.14 or 10.0% (32 d.f.)

SUMMARY OF RESULTS

GRAIN:CWT

	O	C	D	T	Mean
	(1) and (2)				(± 0.66)
O	27.8	32.5	34.0	28.3	30.6
I	30.2	32.1	35.4	31.5	32.3
	NO	N1	N2	N3	
	(1) and (2)				
O	14.9	34.7	39.4	33.5	
I	15.2	36.4	40.7	37.0	
	(3) and (4)				(± 0.64)
O	13.0	32.5	37.8	32.7	29.0
C	15.4	33.3	43.1	37.6	32.3
D	18.3	41.4	42.0	37.2	34.7
T	13.6	35.0	37.4	33.5	29.9
Mean (± 0.64)	15.1	35.6	40.1	35.2	31.5

Mean D.M.%: 84.8

(1) (± 1.03) (3) (± 1.28) For use in vertical and diagonal comparisons only.

(2) (± 0.90) (4) (± 1.28) For use in horizontal and interaction comparisons only.