


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1969

[Full Table of Content](#)



69/R/W/6 - Ccc Treated Seed & N -

Rothamsted Research

Rothamsted Research (1970) *69/R/W/6 - Ccc Treated Seed & N - ; Numerical Results Of The Field Experiments 1969*, pp 247 - 248 - DOI: <https://doi.org/10.23637/ERADOC-1-96>

WINTER WHEAT

(69/R/W/6)

Chlormequat treated seed and N, Long Hoos VII 1969.

Design: 4 randomised blocks of 6 plots.

Area of each plot: 0.0036. Area harvested: 0.0021.

Treatments: All combinations of:-

1. Chlormequat*: None (0), seed treated with chlormequat (C).
2. Nitrogen: 0.5 (N1), 1.0 (N2), 1.5 (N3) cwt as 'Nitro-Chalk' top dressed in spring.

* The amount of chlormequat in the seed was 4.8%.
It was applied by spraying the seed.

Basal applications: 280 lb (0:20:20) broadcast by drill at sowing. Weedkillers: Paraquat at 0.5 lb ion in 25 gals. Mecoprop at 42 oz and 2,4-D at 0.5 oz in 20 gals.

Cultivations, etc.: Ploughed: 6 Aug, 1968. Paraquat applied: 18 Oct. Seed drilled at 190 lb: 22 Oct. 'Nitro-Chalk' applied: 29 Apr, 1969. Mecoprop/2,4-D applied: 3 May. Combine harvested: 1 Sept. Variety: Cappelle. Previous crops: Lucerne 1964 - 1967.

NOTE: A plant count was made on 6 Feb, 1969, a tiller count on 5 June and stem height estimations on 5 and 13 June. Samples for yield were taken on 25 Aug.

Standard error per plot.

Grain, cwt: 4.77 or 9.0% (13 d.f.)

218

SUMMARY OF RESULTS

GRAIN:CWT

	N1	N2	N3	Mean
		(±2.39)		(±1.38)
O	52.2	60.8	59.1	57.4
C	47.2	49.2	48.3	48.2
Mean (±1.69)	49.7	55.0	53.7	52.8

Mean D.M.%: 81.9