


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1969

[Full Table of Content](#)



69/R/W/11 - Gaps - Spring Wheat

Rothamsted Research

Rothamsted Research (1970) *69/R/W/11 - Gaps - Spring Wheat* ; Numerical Results Of The Field Experiments 1969, pp 251 - 252 - DOI: <https://doi.org/10.23637/ERADOC-1-96>

SPRING WHEAT

(69/R/W/11)

Effect of gaps - Great Harpenden II 1969.

Design: 4 randomised blocks of 6 plots.

Area of each plot: 0.0121. Area harvested: 0.0121.

Treatments: No rows missing, 30 rows harvested (G0) and gapping of full plant equivalent to 8 rows removed per plot at growth stages shown below (each plot 17' 6" wide, i.e. nominally 30 rows at 7" spacing, paths 1 ft. wide between plots. The full width was harvested for yield).

At seedling stage:

3 inch gaps evenly spaced (G1)

6 inch gaps evenly spaced (G2)

12 inch gaps evenly spaced (G4)

Pairs of rows removed (R)

At start of stem elongation:

Pairs of rows removed (RL)

Basal applications: 340 lb (25:10:10) combine drilled.

Weedkillers: Paraquat at 0.5 lb ion in 20 gals. 2,4-D at 8 oz and dichlorprop at 32 oz in 20 gals.

Cultivations, etc.: Paraquat applied: 2 Apr, 1969.

Ploughed: 10 Apr. Seed drilled at 160 lb: 16 Apr.

Gapped: 7-14 May. 2,4-D/dichlorprop applied: 21 May.

Gapped: 2 June. Combine harvested: 2 Sept. Variety:

Kolibri. Previous crops: Grass since 1959, Potatoes 1968.

Standard error per plot.

Grain, cwt: 0.57 or 1.2% (15 d.f.)

SUMMARY OF RESULTS

GRAIN: CWT

G0	G1	G2	G4	R	RL	Mean
						(± 0.28)
48.7	45.1	45.3	44.9	43.7	32.6	45.2

Mean D.M. %: 81.8