


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1969

[Full Table of Content](#)



69/R/W/12 - Varieties, N & Ethirimol - Spring Wheat

Rothamsted Research

Rothamsted Research (1970) *69/R/W/12 - Varieties, N & Ethirimol - Spring Wheat* ; Numerical Results Of The Field Experiments 1969, pp 253 - 254 - **DOI:**

<https://doi.org/10.23637/ERADOC-1-96>

SPRING WHEAT

(69/R/W/12)

Varieties, N and ethirimol ('Milstem' or 'PP149'), Whittlocks 1969.

Design: 4 randomised blocks of 10 plots, split into 3.

Area of each sub plot: 0.0080. Area harvested: 0.0053.

Treatments: All combinations of:-

- Whole plots: 1. Varieties: Kloka (KL), Kolibri (KO),
Maris Ensign (ME), Rothwell Sprite (RS),
Troll (T).
2. Fungicide: None (O) ethirimol as seed
dressing at 1 lb (F).
Sub plots: 3. Nitrogen: 0.5 (N1), 1.0 (N2), 1.5 (N3)
cwt (0.5 cwt N in basal compound, re-
mainder as 'Nitro-Chalk').

NOTE: A sixth variety was included but the germination of the seed was poor and the results have been omitted.

Basal applications: 370 lb (15:15:15) combine drilled.
Weedkillers: 2,4-D at 8 oz and dichlorprop at 32 oz
in 20 gals.

Cultivations, etc.: Ploughed: 21 Nov, 1968. Seed drilled
at 185 lb*: 3 Apr, 1969. 'Nitro-Chalk' applied: 11 Apr.
Weedkiller applied: 22 May. Combine harvested: 3 Sept.
Previous crops: Spring beans 1967, winter wheat 1968.

* Rothwell Sprite at 165 lb because its germination
was better than the other varieties.

NOTE: Samples were taken for assessment of mildew
(Erysiphe graminis) and other foliar diseases.

Standard errors per plot.

Grain, cwt: Whole plot: 1.64 or 4.5% (27 d.f.)
Sub plot: 2.16 or 5.9% (60 d.f.)

SUMMARY OF RESULTS

GRAIN:CWT

	KL	KD	ME	RS	T	Mean
	(±0.82)					(±0.37)
O	33.5	42.2	38.4	32.9	36.5	36.7
F	33.0	40.9	37.4	32.9	36.3	36.1
	(1) and (2)					(±0.34)
N1	31.4	37.3	35.9	30.7	35.3	34.1
N2	34.9	43.2	39.6	33.8	38.3	38.0
N3	33.5	44.1	38.1	34.1	35.6	37.1
Mean (±0.58)	33.2	41.5	37.9	32.9	36.4	36.4

Mean D.M.%: 82.6

- (1) (±0.85) For use in horizontal and diagonal comparisons only
 (2) (±0.76) For use in vertical and interaction comparisons only