


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1969

[Full Table of Content](#)



69/R/W/13 - Paths & Blank Rows - Spring Wheat

Rothamsted Research

Rothamsted Research (1970) *69/R/W/13 - Paths & Blank Rows - Spring Wheat* ; Numerical Results Of The Field Experiments 1969, pp 255 - 256 - DOI: <https://doi.org/10.23637/ERADOC-1-96>

SPRING WHEAT

(69/R/W/13)

Effects of paths and blank rows - Whittlocks 1969.

Design: 4 blocks of 8 plots, randomisation restricted.

Area of each plot: 0.0161. Area harvested: 0.0107.

Treatments: All combinations of:-

1. Blank rows: None, 16 middle rows harvested (R0)
2 blank rows, 3 (1) 16 (1) 3 sown (R1)
4 blank rows, 2 (2) 16 (2) 2 sown (R2)
6 blank rows, 1 (3) 16 (3) 1 sown (R3).
2. N: 0.6 (N1), 1.2 (N2) cwt as 'Nitro-Chalk', broadcast by drill.

All plots were 24 row-spaces wide, 16 being harvested. (1) etc. indicate number and position of blank (unsown) rows.

Basal applications: 280 lb (0:20:20) combine drilled. Weedkiller: 2,4-D at 8 oz and dichlorprop at 32 oz in 20 gals.

Cultivations, etc.: Ploughed: 21 Nov, 1968. Seed drilled at 180 lb per sown acre: 2 Apr, 1969. 'Nitro-Chalk' applied: 10 Apr. Weedkiller applied: 22 May. Combine harvested: 3 Sept. Variety: Kolibri. Previous crops: Spring beans 1967, winter wheat 1968.

Standard error per plot.

Grain, cwt: 1.92 or 4.4% (21 d.f.)

SUMMARY OF RESULTS

GRAIN: CWT

	R0	R1	R2	R3	Mean
	(±0.96)				(±0.48)
N1	38.4	39.6	43.4	43.3	41.2
N2	43.7	45.0	46.2	46.2	45.3
Mean (±0.68)	41.0	42.3	44.8	44.7	43.2

Mean D.M.%: 84.2