


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1969

[Full Table of Content](#)



69/R/B/4 - Early & Late Mildew - Barley

Rothamsted Research

Rothamsted Research (1970) *69/R/B/4 - Early & Late Mildew - Barley* ; Numerical Results Of The Field Experiments 1969, pp 265 - 266 - DOI: <https://doi.org/10.23637/ERADOC-1-96>

BARLEY

(69/R/B/4)

Early and late mildew (*Erysiphe graminis*), Long Hoos I and II 1969.

Design: 4 randomised blocks of 4 plots.

Area of each plot: 0.0321. Area harvested: 0.0129.

Treatments: Fungicide ethirimol ('PP149' or 'Milstem'):-

None

(0)

0.25 lb as seed dressing

(1D)

2 lb as seed dressing, crop sprayed on 2 occasions, the first at 1 lb and the second at 1.5 lb in 30 gals.

(8D)

Crop sprayed on 2 occasions as above

(4S)

Basal applications: 340 lb (20:10:10) combine drilled.

Weedkiller: Mecoprop at 36 oz and 2,4-D at 9 oz in 20 gals.

Cultivations, etc.: Deep-tine cultivated 3 times: 13 Nov, 1968.

Seed drilled at 140 lb: 27 Mar, 1969. Weedkiller applied:

15 May. Ethirimol spray applied: 25 June and 10 July. Combine harvested: 14 Aug. Variety: Zephyr. Previous crops:

Fallow 1967, potatoes 1968.

NOTE: Samples were taken for assessment of mildew (*Erysiphe graminis*) and other foliar diseases and for assessment of tiller number, tiller length and ear sizes.

Standard error per plot.

Grain, cwt: 2.09 or 4.2% (9 d.f.)

SUMMARY OF RESULTS

0	1D	8D	4S	Mean
GRAIN: CWT				
(±1.05)				
48.4	48.6	52.6	49.1	49.7
STRAW: CWT				
32.5	31.3	36.1	34.0	33.5

Mean D.M.%: Grain: 80.5
 Straw: 82.7