

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1970

[Full Table of Content](#)



70/R/CS/13 N and Mg Levels to Old Grass

Rothamsted Research

Rothamsted Research (1971) *70/R/CS/13 N and Mg Levels to Old Grass* ; Numerical Results Of The Field Experiments 1970, pp 143 - 146 - DOI: <https://doi.org/10.23637/ERADOC-1-59>

N AND Mg LEVELS TO OLD GRASS

(70/R/CS/13)

Old grass, Park Grass plot 6, 1970, the sixth year.

For previous years' results see 'Results' 65/C/33, 66/C/14, 67/C/10, 68/C/8 and 69/R/CS/13.

Area of each plot: 0.0045. Area harvested: 0.0021.

Additional treatments: A test of 25 (M1) v 50 lb (M2) Mg as magnesium sulphate applied in spring has been added in combination with the other residual treatments. A certain high order interaction is confounded with block differences. All plots are now cut 4 times a year and the 'Nitro-Chalk' is applied in equal dressings for each cut. Total N rates per year are unchanged. Basal applications of sodium and magnesium have been discontinued.

Basal application: 23 cwt ground chalk.

Cultivations, etc.: Basal PK applied: 20 Feb, 1970. Test Mg applied: 23 Feb. N applied: 16 Mar. Ground chalk applied: 16 Apr. S plots sprayed with mecoprop at 45 oz a.e. in 40 gals: 1 May. S plots spot-sprayed with mecoprop at the same rate: 1 Sept. Plots cut: 20 May, 16 July, 4 Sept, 23 Oct. N applied after each cut except the last.

Standard errors per plot. Dry matter, cwt:

1st cut:	1.78 or 7.8% (14 d.f.)
2nd cut:	2.10 or 19.6% (14 d.f.)
3rd cut:	1.71 or 10.7% (14 d.f.)
4th cut:	1.16 or 13.9% (14 d.f.)
Total of all 4 cuts:	3.81 or 6.6% (14 d.f.)

70/R/CS/13

SUMMARY OF RESULTS

DRY MATTER: CWT

	NO	N1	N2	N3	Mean	SNO
1ST CUT						
1969					(±0.45)	
		(±0.89)				
C3	10.4	15.7	29.6	35.7	22.8	1.9
C6	9.4	16.6	30.1	35.7	22.9	1.7
1970						
M1	9.7	15.9	29.8	35.7	22.8	1.8
M2	10.1	16.3	29.8	35.7	23.0	1.8
Mean (±0.63)	9.9	16.1	29.8	35.7	22.9	1.8
2ND CUT						
1969					(±0.53)	
		(±1.05)				
C3	12.7	11.6	12.1	7.5	11.0	3.9
C6	11.4	11.6	11.3	7.5	10.4	3.5
1970						
M1	12.7	11.6	12.4	8.0	11.2	3.7
M2	11.4	11.6	11.0	6.9	10.2	3.7
Mean (±0.74)	12.0	11.6	11.7	7.5	10.7	3.7
1st cut:	General mean:		Mean D.M. %:			
2nd cut:	18.7		19.6			
	9.3		28.0			

70/R/CS/13

DRY MATTER: CWT

	NO	N1	N2	N3	Mean	SNO
3RD CUT						
1969	(±0.85)				(±0.43)	
C3	11.1	12.2	20.2	19.9	15.9	3.2
C6	9.9	12.9	19.5	21.6	16.0	3.8
1970						
M1	10.5	13.0	20.4	21.1	16.3	3.2
M2	10.4	12.1	19.3	20.4	15.6	3.8
Mean (±0.60)	10.5	12.6	19.9	20.8	15.9	3.5
4TH CUT						
1969	(±0.58)				(±0.29)	
C3	4.1	6.7	11.6	10.7	8.3	3.0
C6	4.1	6.5	11.0	12.0	8.4	2.2
1970						
M1	4.4	7.2	11.4	11.5	8.6	2.7
M2	3.8	6.0	11.3	11.2	8.1	2.4
Mean (±0.41)	4.1	6.6	11.3	11.4	8.4	2.6
General mean: Mean D.M. %:						
3rd cut:	13.4			22.8		
4th cut:	7.2			25.0		

70/R/CS/13

DRY MATTER: CWT

	NO	N1	N2	N3	Mean	SNO
TOTAL OF 4 CUTS						
1969					(±0.95)	
		(±1.90)				
C3	38.2	46.3	73.6	73.8	58.0	12.0
C6	34.8	47.6	71.9	76.9	57.8	11.1
1970						
M1	37.3	47.8	74.1	76.4	58.9	11.5
M2	35.7	46.1	71.5	74.3	56.9	11.7
Mean (±1.35)	36.5	47.0	72.8	75.3	57.9	11.6

General mean: 48.6

Mean D.M. %: 23.8