

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1970

[Full Table of Content](#)



70/W/CS/55 Spring Beans

Rothamsted Research

Rothamsted Research (1971) *70/W/CS/55 Spring Beans ; Numerical Results Of The Field Experiments 1970*, pp 201 - 202 - DOI: <https://doi.org/10.23637/ERADOC-1-59>

SPRING BEANS

(70/W/CS/55)

Effects of fumigation and nitrogen - Woburn Butt Furlong 1970, second year. For previous year's results see 'Results' 69/W/BE/1.

Design: 3 blocks of 6 plots.

Area of each plot: 0.0022. Area harvested: 0.0005.

Treatments (cumulative 1969, 1970): All combinations of:-

1. Fumigant: None (0), dazomet at 400 lb (F).
2. Nitrogen: None (N0), 1.0 (N1), 2.0 (N2) cwt N as 'Nitro-Chalk' 21 half seedbed, half top dressed.

Basal applications: 400 lb (0:14:28). Insecticide: Demeton-s-methyl at 3.5 fl oz in 35 gals.

Cultivations, etc.: Ploughed: 9 Sept, 1969. Fumigant applied, rotary cultivated twice to 4 inches and 6 inches: 18 Sept. Ploughed second time: 11 Feb, 1970. Power harrowed, fertiliser placed, seed drilled at 200 lb: 20 Mar. First application of N: 3 Apr. Second application of N: 15 May. Insecticide applied: 22 June. Hand harvested: 28 Aug. Variety: Tarvin.

- NOTES: (1) Soil samples were taken for counts of ectoparasitic nematodes.
- (2) Plant samples were taken for observations of fungal pathogens.
- (3) Counts were made of germination, number of stems, number of pods, 1000 grain weights and percentage nitrogen in grain.

Standard error per plot.

Grain, cwt: 0.72 or 7.3% (10 d.f.)

70/W/CS/55

SUMMARY OF RESULTS

GRAIN: CWT

	NO	N1	N2	Mean
		(± 0.41)		(± 0.24)
O	9.6	9.5	5.5	8.2
F	9.7	11.5	12.9	11.4
Mean (± 0.29)	9.6	10.5	9.2	9.8

Mean D.M. %: 83.0