

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1970

[Full Table of Content](#)



70/R/AG/6 Agdell - Grass and Fallow, Sugar Beet and Barley

Rothamsted Research

Rothamsted Research (1971) *70/R/AG/6 Agdell - Grass and Fallow, Sugar Beet and Barley ; Numerical Results Of The Field Experiments 1970*, pp 24 - 34 - **DOI:**
<https://doi.org/10.23637/ERADOC-1-59>

AGDELL

(70/R/AG/6)

Grass, sugar beet and barley 1970.

First year of revised scheme testing fresh P on two phases per annum of a three course rotation - sugar beet, barley, potatoes. Crops in 1970 sugar beet and barley.

For history, past treatments etc. and for previous years' results. see 'Details' 1967 and 'Results' 68/A/4 and 69/R/AG/6.

Area of each sub-plot:

Grass and fallow: Plots 1, 2, 3, 4 - 0.0180. Plots 5, 6 - 0.0162.

Area harvested: Plots 1, 2, 3, 4 - 0.0023. Plots 5, 6 - 0.0020.

Sugar beet and barley: Plots 1, 2, 3, 4 - 0.0045. Plots 5, 6 - 0.0040.

Area harvested: Sugar beet: Plots 1, 2, 3, 4 - 0.0019. Plots 5, 6 - 0.0017. Barley: Plots 1, 2, 3, 4 - 0.0017. Plots 5, 6 - 0.0015.

New superphosphate treatments:

Sugar beet: None (P0), 1.0 cwt (P2) P205

Barley: None (P0), 0.5 cwt (P1) P205
(P as single superphosphate).

Only the sub-plots formerly testing P have been brought into the new scheme. Pairs of these sub-plots were split in a North-South direction for crops. Each original sub-plot was further split in an East-West direction for the test of fresh superphosphate.

Treatments and cropping on sub-plots testing K were unaltered.

Balancing P and K: P (as triple superphosphate) and K (as muriate of potash) were applied on 18 Nov, 1969, to balance removals by grass in 1969, to all sub-plots except P0, which continues to receive no P, and K0, which continues to receive no K.

70/R/AG/6

Plot No	Rates in cwt P205				Sub-plots testing K:-			
	Sub-plots testing P:-							
	P0	P1	P2	P4	K0	K1	K2	K4
1	0	0.41	0.44	0.47	0.35	0.45	0.46	0.49
2	0	0.36	0.43	0.47	0.31	0.44	0.52	0.47
3	0	0.35	0.39	0.39	0.34	0.39	0.39	0.39
4	0	0.40	0.36	0.45	0.42	0.44	0.40	0.46
5	0	0.29	0.34	0.39	0.29	0.36	0.36	0.35
6	0	0.31	0.36	0.36	0.22	0.42	0.41	0.44

	Rates in cwt K20				Sub-plots testing K:-			
	Sub-plots testing P:-							
	P0	P1	P2	P4	K0	K1	K2	K4
1	2.59	3.00	3.13	2.92	0	2.36	2.60	3.08
2	2.24	2.67	2.93	2.99	0	2.20	2.90	2.81
3	1.99	2.47	2.75	2.44	0	2.38	2.18	2.53
4	1.30	2.80	2.36	2.84	0	2.46	2.19	2.77
5	0.99	2.25	2.26	2.48	0	1.96	2.10	2.23
6	0.81	2.20	2.26	2.21	0	2.26	2.20	2.45

Basal applications: Manures:-

Sugar beet: 1.5 cwt N as 'Nitro-Chalk', 2.5 cwt K2O as muriate of potash, 0.8 cwt MgO as kieserite.

Barley: 0.75 cwt N, 0.48 cwt K2O as (25:0:16).

Grass: as previously.

Insecticide to sugar beet: Menazon ('Saphicol' at 0.5 pints in 40 gals) on 2 occasions.

Cultivations, etc.: Ground chalk applied at 25 cwt to main plots 1 and 2 and to South half of 3 and 4: 31 Oct, 1969.

Grass: Basal N applied: 18 Mar, 1970 and 15 June. Cut twice for silage: 4 June and 3 Aug. Ploughed: 4 Sept.

Fallow: Ploughed: 5 Dec, 1969. Rotary cultivated twice: 10 June, 1970 and 22 July.

Sugar beet: Ploughed: 5 Dec, 1969. Test P and basal NK Mg applied: 30 Apr, 1970. Seed drilled at 8 lb: 4 May. Singled: 8 June.

Insecticide applied: 25 June, 15 July. Lifted: 19 Oct.

Variety: Klein E.

Barley: Ploughed: 5 Dec, 1969. Test P applied, seed drilled at 150 lb, basal N and K applied: 24 Apr, 1970. Combine harvested: 18 Aug. Variety: Julia.

Errata to 'Results' 1969: Delete table on p.24 of 'Rates in cwt K20' and replace by:

70/R/AG/6

Plot No	Sub-plots testing P:-				Sub-plots testing K:-			
	P0	P1	P2	P4	K0	K1	K2	K4
1	2.23	3.31	2.72	3.23	0	2.64	3.02	3.06
2	1.40	2.85	3.25	3.08	0	2.66	2.89	2.88
3	1.32	2.66	2.77	2.97	0	2.54	2.38	2.72
4	0.89	2.88	2.66	2.67	0	2.85	2.76	3.09
5	0.65	2.26	2.46	2.67	0	2.19	2.54	2.80
6	0.68	2.42	2.56	2.61	0	2.42	2.52	2.49

70/R/AG/6

SUMMARY OF RESULTS

GRASS

DRY MATTER: CWT

Plot

K	5	6	3	4	1	2	Mean
1ST CUT							
0	14.5	16.2	25.0	22.8	27.5	17.4	20.6
1	36.2	40.6	36.9	39.2	40.8	42.7	39.4
2	36.4	45.2	40.5	41.9	39.1	44.5	41.3
4	38.6	40.7	41.4	42.0	39.2	45.1	41.2
Mean	31.4	35.7	36.0	36.5	36.7	37.4	35.6
2ND CUT							
0	3.6	2.6	3.1	3.6	4.4	1.6	3.2
1	10.9	13.2	8.9	14.1	11.1	11.9	11.7
2	9.5	10.7	9.2	11.3	12.3	22.1	12.5
4	8.6	10.3	8.9	13.6	10.6	14.3	11.0
Mean	8.2	9.2	7.5	10.6	9.6	12.5	9.6
TOTAL OF 2 CUTS							
0	18.1	18.8	28.1	26.4	31.8	19.1	23.7
1	47.0	53.8	45.8	53.3	52.0	54.6	51.1
2	45.9	55.9	49.7	53.2	51.4	66.5	53.8
4	47.2	51.0	50.3	55.5	49.8	59.4	52.2
Mean	39.6	44.9	43.5	47.1	46.3	49.9	45.2
Mean D.M. %:							
1st cut:			27.1				
2nd cut:			28.9				
Total of 2 cuts:			28.0				

70/R/AG/6

SUGAR BEET

ROOTS (WASHED): TONS

		Plot						Mean
Treatment 1848-1957		5 None	6	3 PK	4	1 NPK	2	
P	Previous cropping 1958-69, Arable or fallow							
1964-69 1970								
0	0	8.15	13.63	15.90	11.22	12.51	13.21	12.44
1	0	11.09	13.36	14.03	11.34	15.90	11.46	12.86
2	0	13.76	13.50	16.37	13.68	12.39	16.72	14.40
4	0	15.23	14.83	12.98	16.49	16.14	15.78	15.24
0	2	12.03	14.83	16.84	15.90	15.78	12.98	14.73
1	2	13.50	9.49	13.91	10.99	15.55	12.51	12.66
2	2	16.97	11.49	17.89	14.62	14.97	16.49	15.40
4	2	11.76	13.63	11.81	14.50	17.89	12.74	13.72
Mean		12.81	13.10	14.97	13.59	15.14	13.99	13.93
P	Previous cropping 1958-69, Grass							
1964-69 1970								
0	0	6.81	1.07	4.79	8.54	12.74	12.28	7.71
1	0	14.57	16.17	16.84	16.95	17.54	16.72	16.46
2	0	18.04	13.76	15.67	16.60	13.68	19.06	16.14
4	0	10.56	17.77	16.60	12.39	16.49	17.77	15.26
0	2	14.03	11.22	11.46	12.16	15.55	15.08	13.25
1	2	13.50	15.90	16.95	15.32	17.77	17.42	16.14
2	2	18.57	13.63	16.02	13.68	13.33	16.14	15.23
4	2	15.63	17.50	14.97	13.91	14.26	16.49	15.46
Mean		13.96	13.38	14.16	13.69	15.17	16.37	14.46

70/R/AG/6

SUGAR BEET

SUGAR %

Plot

Treatment 1848-1957		5 None	6	3 PK	4	1 NPK	2	Mean
P	Previous cropping 1958-69, Arable or fallow							
1964-69	1970							
0	0	16.3	16.7	17.6	14.8	16.9	16.1	16.4
1	0	17.1	16.6	17.2	15.2	17.2	16.0	16.5
2	0	16.0	16.4	16.7	16.8	16.4	16.7	16.5
4	0	17.2	17.2	17.0	15.9	17.0	17.2	16.9
0	2	15.3	17.0	16.9	16.0	16.5	16.0	16.3
1	2	16.3	16.3	16.8	16.2	16.7	15.9	16.4
2	2	16.5	16.1	16.8	16.6	16.6	16.6	16.5
4	2	15.6	16.8	17.1	16.6	16.2	16.4	16.5
Mean		16.3	16.6	17.0	16.0	16.7	16.3	16.5
P	Previous cropping 1958-69, Grass							
1964-69	1970							
0	0	16.0	16.6	14.9	15.2	16.3	16.0	15.8
1	0	17.2	16.7	17.4	16.3	16.8	16.4	16.8
2	0	16.5	17.0	17.4	16.3	16.7	16.8	16.8
4	0	17.0	16.5	17.0	16.4	16.8	17.1	16.8
0	2	16.4	15.9	16.1	15.7	16.4	16.6	16.2
1	2	16.5	16.5	17.3	16.8	16.7	17.3	16.9
2	2	17.3	16.9	17.2	16.5	16.1	16.7	16.8
4	2	16.9	17.0	17.7	16.4	16.8	17.1	17.0
Mean		16.7	16.7	16.9	16.2	16.6	16.8	16.6

70/R/AG/6

SUGAR BEET

TOTAL SUGAR: CWT

Plot

Treatment 1848-1957		5 None	6	3	4 PK	1	2 NPK	Mean
P	Previous cropping 1958-69, Arable or fallow							
1964-69 1970								
0	0	26.5	45.5	56.1	33.2	42.3	42.6	41.0
1	0	37.8	44.3	48.3	34.5	54.8	36.6	42.7
2	0	44.1	44.3	54.7	45.9	40.7	55.8	47.6
4	0	52.3	50.9	44.1	52.5	54.7	54.2	51.5
0	2	36.7	50.4	56.8	50.7	52.1	41.5	48.1
1	2	43.9	30.9	46.8	35.5	52.0	39.7	41.5
2	2	56.1	36.9	60.1	48.4	49.7	54.6	51.0
4	2	36.7	45.8	40.5	48.1	58.0	41.8	45.1
Mean		41.8	43.6	50.9	43.6	50.5	45.9	46.1
P	Previous cropping 1958-69, Grass							
1964-69 1970								
0	0	21.9	3.5	14.3	26.0	41.5	39.3	24.4
1	0	50.2	54.1	58.6	55.3	59.1	54.9	55.4
2	0	59.7	46.9	54.7	54.0	45.7	64.2	54.2
4	0	36.0	58.8	56.3	40.6	55.3	60.9	51.3
0	2	46.0	35.7	36.9	38.1	51.1	50.1	43.0
1	2	44.5	52.6	58.8	51.4	59.3	60.4	54.5
2	2	64.2	46.0	55.0	45.0	42.9	53.8	51.1
4	2	52.9	59.6	52.9	45.7	48.0	56.4	52.6
Mean		46.9	44.7	48.4	44.5	50.3	55.0	48.3

70/R/AG/6

SUGAR BEET

TOPS: TONS

		Plot						
Treatment 1848-1957		5 None	6	3	4 PK	1	2 NPK	Mean
P		Previous cropping 1958-69, Arable or fallow						
1964-69	1970							
0	0	12.98	18.24	21.75	15.90	19.18	19.41	17.91
1	0	11.34	17.07	19.64	16.84	21.28	16.37	17.09
2	0	13.45	18.94	19.18	16.37	17.77	21.28	17.83
4	0	18.82	17.19	16.60	18.01	21.51	18.94	18.51
0	2	16.25	18.47	21.28	23.38	22.68	23.38	20.91
1	2	14.73	16.14	20.11	16.84	23.62	19.64	18.51
2	2	16.25	16.60	21.98	22.68	19.18	21.51	19.70
4	2	15.55	17.19	19.41	20.11	22.45	18.47	18.86
Mean		14.92	17.48	19.99	18.77	20.96	19.88	18.67
P		Previous cropping 1958-69, Grass						
1964-69	1970							
0	0	7.02	1.14	6.78	9.59	18.24	15.90	9.78
1	0	17.54	19.99	25.72	20.11	24.09	24.09	21.92
2	0	18.59	17.77	20.81	20.58	22.45	26.42	21.10
4	0	11.93	20.23	22.68	16.60	23.62	27.36	20.40
0	2	19.29	16.02	21.28	16.60	24.32	23.85	20.23
1	2	13.33	22.22	23.62	20.11	25.49	23.38	21.36
2	2	19.41	14.26	22.22	18.47	20.11	25.49	19.99
4	2	17.77	20.93	18.24	16.37	21.98	23.62	19.82
Mean		15.61	16.57	20.17	17.30	22.54	23.76	19.33

70/R/AG/6

SUGAR BEET

PLANT NUMBER: THOUSANDS

		Plot						Mean
Treatment 1848-1957		5 None	6	3 PK	4	1 NPK	2	
P		Previous cropping 1958-69, Arable or fallow						
1964-69	1970							
0	0	16.8	19.4	26.2	17.8	30.9	25.1	22.7
1	0	15.2	22.5	27.2	16.2	31.4	23.0	22.6
2	0	14.1	18.3	24.6	18.9	25.7	25.1	21.1
4	0	22.5	19.4	23.6	15.7	27.8	24.6	22.3
0	2	18.3	17.8	25.7	28.3	33.0	24.1	24.5
1	2	18.3	17.3	21.0	16.8	30.9	24.1	21.4
2	2	15.2	15.2	24.6	20.4	29.3	24.6	21.6
4	2	14.1	14.1	23.0	19.9	26.7	22.5	20.1
Mean		16.8	18.0	24.5	19.3	29.5	24.2	22.0
P		Previous cropping 1958-69, Grass						
1964-69	1970							
0	0	22.5	18.6	23.6	18.3	28.3	26.7	23.0
1	0	16.8	21.0	32.0	21.0	23.0	26.2	23.3
2	0	21.0	16.2	22.5	22.0	28.3	26.7	22.8
4	0	13.1	20.4	21.0	16.8	29.3	29.9	21.7
0	2	24.1	18.9	21.5	17.8	29.3	27.8	23.2
1	2	13.1	21.0	23.6	20.4	28.3	27.8	22.3
2	2	25.1	17.3	23.0	16.8	28.8	24.6	22.6
4	2	17.3	20.4	19.9	19.4	24.6	30.4	22.0
Mean		19.1	19.2	23.4	19.1	27.5	27.5	22.6

70/R/AG/6

BARLEY

GRAIN: CWT

Plot

Treatment 1848-1957		5 None	6	3	4 PK	1	2 NPK	Mean
P		Previous cropping 1958-69, Arable or fallow						
1964-69	1970							
0	0	9.1	13.8	34.3	14.9	14.7	19.9	17.8
1	0	23.1	21.5	36.7	25.3	21.8	11.1	23.3
2	0	29.4	37.8	37.5	40.0	20.7	24.6	31.7
4	0	29.6	39.3	34.8	43.0	25.7	21.7	32.4
0	1	8.0	25.1	33.4	14.7	18.2	15.0	19.1
1	1	28.2	27.1	36.9	30.5	14.9	14.3	25.3
2	1	36.3	36.2	35.3	41.0	21.1	31.0	33.5
4	1	32.0	38.5	39.3	37.7	29.6	23.3	33.4
Mean		24.5	29.9	36.0	30.9	20.8	20.1	27.0

P		Previous cropping 1958-69, Grass						
1964-69	1970							
0	0	18.8	16.6	28.4	18.2	26.9	24.5	22.2
1	0	33.2	36.0	36.3	29.2	35.1	30.5	33.4
2	0	32.5	41.5	35.7	26.0	25.7	29.2	31.8
4	0	34.7	37.6	37.6	42.6	30.1	34.6	36.2
0	1	17.9	17.1	30.0	19.0	29.5	29.3	23.8
1	1	28.4	38.8	37.5	31.6	28.5	30.0	32.5
2	1	32.1	37.6	35.4	32.8	26.8	32.1	32.8
4	1	38.3	38.3	39.7	25.8	34.3	34.7	35.2
Mean		29.5	32.9	35.1	28.2	29.6	30.6	31.0

Mean D.M. %: 84.7

70/R/AG/6

BARLEY

STRAW: CWT

Plot

Treatment 1848-1957		5 None	6	3 PK	4	1 NPK	2	Mean
P	Previous cropping 1958-69, Arable or fallow							
1964-69 1970								
0	0	4.3	10.4	19.2	9.8	10.8	14.5	11.5
1	0	13.9	13.4	24.5	15.8	13.8	7.0	14.7
2	0	16.1	21.8	21.2	23.3	13.7	16.0	18.7
4	0	14.5	20.7	19.6	24.2	16.2	12.9	18.0
0	1	3.1	12.6	19.0	8.6	16.8	11.0	11.9
1	1	16.5	16.0	24.2	19.7	18.1	8.9	17.2
2	1	21.7	20.6	19.0	23.0	13.1	20.3	19.6
4	1	14.9	20.4	23.8	23.8	16.4	11.3	18.4
Mean		13.1	17.0	21.3	18.5	14.9	12.7	16.3

P	Previous cropping 1958-69, Grass							
1964-69 1970								
0	0	7.6	9.3	15.1	8.8	17.7	14.3	12.1
1	0	16.2	17.7	19.2	16.9	19.5	20.5	18.3
2	0	16.2	19.9	20.4	14.9	22.5	20.1	19.0
4	0	18.6	20.1	23.2	24.1	20.4	22.5	21.5
0	1	8.6	8.2	14.8	11.0	17.6	16.6	12.8
1	1	14.8	19.7	21.9	18.6	17.2	23.5	19.3
2	1	14.5	19.0	19.6	19.7	20.1	20.7	18.9
4	1	19.8	19.6	22.4	22.8	22.2	20.6	21.2
Mean		14.5	16.7	19.6	17.1	19.6	19.9	17.9

Mean D.M. %: 54.0