


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1971

[Full Table of Content](#)



71/W/RN/14 Long-term Phosphate - Barley, Potatoes

Rothamsted Research

Rothamsted Research (1972) *71/W/RN/14 Long-term Phosphate - Barley, Potatoes* ; Yields Of The Field Experiments 1971, pp 116 - 119 - DOI: <https://doi.org/10.23637/ERADOC-1-97>

71/W/RN/14

LONG TERM PHOSPHATE

Object: To study direct and residual effects of superphosphate on yields of three crops grown in rotation - Woburn Stackyard III.

The fourth year, barley and potatoes.

For previous years see 68/B/8(t), 69/W/RN/14 and 70/W/RN/14(t).

Design: 6 blocks of 6 plots, split into 2.

Whole plot dimensions: 8.53 x 15.8. Area harvested: Barley - 0.00442, potatoes - 0.00451.

Treatments:

Sub plots: Superphosphate:

Barley: None (0), 63 kg P₂O₅ (P).

Potatoes: None (0), 188 kg P₂O₅ (P).

Basal applications:-

Barley: Manures: 2510 kg ground chalk, 61 kg of K₂O as muriate of potash, 148 kg N as 'Nitro-Chalk' 21. Weedkiller: Ioxynil at 0.525 kg and mecoprop at 1.58 kg in 281 l.

Potatoes: Manures: 2510 kg ground chalk, 258 kg of K₂O as muriate of potash in autumn, 250 kg of K₂O as muriate of potash in spring, 251 kg N as 'Nitro-Chalk' 21, 99 kg MgO as Epsom salts. Weedkiller: Linuron at 1.12 kg in 371 l. Fungicide and insecticide: Mancozeb at 1.34 kg and demeton-s-methyl at 246 ml in 416 l. Fungicide: Mancozeb at 1.34 kg in 371 l.

Cultivations, etc.:

Barley: Ground chalk applied: 26 Nov, 1970. Ploughed: 10 Dec. N and K applied: 1 Mar, 1971. P applied: 4 Mar. Seed drilled at 157 kg: 23 Mar. Power harrowed, redrilled because of bird damage: 13 Apr. Sprayed weedkiller: 18 May. Combine harvested: 25 Aug. Variety: Julia.

Potatoes: Sub soiled, tines 142 cm apart and 61 cm deep, ground chalk and K applied: 11 Sept, 1970. Deep-tine cultivated: 12 Sept. Ploughed: 21 Oct. P applied: 29 Mar, 1971. N, K and Mg applied: 1 Apr. Rotary cultivated, potatoes planted: 2 Apr. Weedkiller applied: 5 May. Rotary ridged: 1 June. Fungicide with insecticide applied: 28 June. Fungicide applied: 12 Aug. Haulm destroyed mechanically: 13 Sept. Lifted: 14 Sept. Variety: Pentland Crown.

71/W/RN/14

Standard errors per plot.

Barley: Grain, tonnes/hectare: Whole plot: 0.234 or 6.0% (9 d.f.)

Sub plot: 0.343 or 8.8% (10 d.f.)

Potatoes: Total tubers, tonnes/hectare:

Whole plot: 2.71 or 7.1% (10 d.f.)

Sub plot: 3.88 or 10.1% (12 d.f.)

71/W/RN/14

SUMMARY OF RESULTS

BARLEY

	RO*	R1	R2	R4	R6	Mean
GRAIN: TONNES/HECTARE						
	(1) & (2)	(3) & (4)				(±0.081)
O	3.15	3.67	3.91	4.08	4.21	3.69
P	3.81	4.31	3.98	4.45	4.37	4.12
Mean	3.48 (±0.096)	3.99	3.94 (±0.135)	4.26	4.29	3.91

(1) (±0.138) (3) (±0.194) For use in horizontal and diagonal comparisons only
 (2) (±0.140) (4) (±0.198) For use in vertical and interaction comparisons only

STRAW: TONNES/HECTARE

O	2.11	2.65	2.87	3.04	2.87	2.61
P	2.76	2.98	2.83	3.10	3.15	2.93
Mean	2.43	2.81	2.85	3.07	3.01	2.77

Mean D.M. %: Grain: 80.6
 Straw: 83.5

* Duplicated treatment

71/W/RN/14

POTATOES

	RO*	R1	R2	R4	R6	Mean
TOTAL TUBERS: TONNES/HECTARE						
	(1) & (2)	(3) & (4)				(±0.91)
O	32.1	34.4	37.0	41.8	38.4	36.0
P	38.5	42.0	43.2	41.7	40.1	40.7
Mean	35.3 (±1.11)	38.2	40.1 (±1.56)	41.8	39.2	38.3

- (1) (±1.57) (3) (±2.23) For use in horizontal and diagonal comparisons only
 (2) (±1.58) (4) (±2.24) For use in vertical and interaction comparisons only

WARE: 3.81 CM (1.5 INCH) RIDDLE

O	97.7	97.2	97.8	98.0	97.2	97.6
P	97.1	96.8	97.7	97.4	97.0	97.2
Mean	97.4	97.0	97.7	97.7	97.1	97.4

* Duplicated treatment