


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1971

[Full Table of Content](#)



71/R/CS/13 N and Mg Levels to Old Grass - Old Grass

Rothamsted Research

Rothamsted Research (1972) *71/R/CS/13 N and Mg Levels to Old Grass - Old Grass* ; Yields Of The Field Experiments 1971, pp 139 - 142 - DOI: <https://doi.org/10.23637/ERADOC-1-97>

71/R/CS/13

N AND Mg LEVELS TO OLD GRASS

Object: To study the effects of a range of nitrogen levels on the production of very old permanent pasture. To see whether application of magnesium sulphate will increase the magnesium content of such pasture when large dressings of nitrogen are applied - Park Grass old plot 6.

The seventh year, old grass.

For previous years see 65/C/33(t), 66/C/14, 67/C/10(t), 68/C/8(t) and 69-70/R/CS/13(t).

Whole plot dimensions: 1.83 x 10.1. Area harvested: 0.00086.

Cultivations, etc.: Basal PK and test Mg applied: 8 Dec, 1970. N applied: 9 Mar, 1971. S plots sprayed with mecoprop at 3.14 kg a.e. in 449 1: 22 Apr and 12 July. Plots cut: 13 May, 21 June, 2 Aug, 14 Oct. N applied after each cut except the last.

Standard errors per plot. Dry matter, tonnes/hectare:

1st cut:	0.358 or 12.3% (14 d.f.)
2nd cut:	0.205 or 8.6% (14 d.f.)
3rd cut:	0.219 or 10.7% (14 d.f.)
4th cut:	0.182 or 6.9% (14 d.f.)
Total of 4 cuts:	0.486 or 4.9% (14 d.f.)

71/R/CS/13

SUMMARY OF RESULTS

DRY MATTER: TONNES/HECTARE

1971

	NO	N1	N2	N3	Mean	SNO
1ST CUT						
1969	(±0.179)				(±0.089)	
C3	1.08	2.21	4.02	4.46	2.95	0.45
C6	1.15	2.23	3.94	4.16	2.87	0.44
1971						
M1	1.12	2.24	3.89	4.28	2.88	0.42
M2	1.11	2.20	4.07	4.34	2.93	0.47
Mean (±0.126)	1.12	2.22	3.98	4.31	2.91	0.44

2ND CUT

1969	(±0.103)				(±0.051)	
C3	1.84	2.53	2.69	2.46	2.38	1.01
C6	1.79	2.44	2.73	2.52	2.37	0.92
1971						
M1	1.76	2.58	2.71	2.49	2.39	1.00
M2	1.87	2.40	2.70	2.49	2.37	0.93
Mean (±0.073)	1.82	2.49	2.71	2.49	2.38	0.96

	General mean:	Mean D.M. %:
1st cut:	2.41	21.2
2nd cut:	2.09	17.8

71/R/CS/13

DRY MATTER: TONNES/HECTARE

1971

	N0	N1	N2	N3	Mean	SNO
3RD CUT						
1969	(±0.109)				(±0.055)	
C3	1.66	1.67	2.34	2.28	1.99	0.51
C6	1.88	1.63	2.25	2.61	2.09	0.68
1971						
M1	1.78	1.66	2.29	2.49	2.05	0.48
M2	1.75	1.65	2.30	2.40	2.03	0.70
Mean (±0.077)	1.77	1.65	2.30	2.44	2.04	0.59

4TH CUT

1969	(±0.091)				(±0.045)	
C3	1.83	2.30	2.80	3.28	2.55	0.90
C6	1.83	2.39	3.07	3.45	2.69	0.89
1971						
M1	1.81	2.42	2.90	3.32	2.61	0.90
M2	1.85	2.28	2.96	3.42	2.63	0.90
Mean (±0.064)	1.83	2.35	2.93	3.37	2.62	0.90

	General mean:	Mean D.M. %:
3rd cut:	1.75	21.9
4th cut:	2.27	21.4

71/R/CS/13

DRY MATTER: TONNES/HECTARE

1971

	NO	N1	N2	N3	Mean	SNO
	TOTAL OF 4 CUTS					
1969	(±0.243)				(±0.121)	
C3	6.41	8.71	11.85	12.49	9.86	2.87
C6	6.65	8.70	11.99	12.74	10.02	2.92
1971						
M1	6.47	8.89	11.80	12.58	9.93	2.78
M2	6.58	8.52	12.04	12.65	9.95	3.00
Mean (±0.172)	6.53	8.70	11.92	12.61	9.94	2.89

General mean: 8.53

Mean D.M. %: 20.6