


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1971

[Full Table of Content](#)



71/S/CS/1 Row Spacing and Seed Rates After Intensive Wheat - Wheat

Rothamsted Research

Rothamsted Research (1972) *71/S/CS/1 Row Spacing and Seed Rates After Intensive Wheat - Wheat* ; Yields Of The Field Experiments 1971, pp 233 - 234 - **DOI:**
<https://doi.org/10.23637/ERADOC-1-97>

71/S/CS/1

ROW SPACING AND SEED RATES AFTER INTENSIVE WHEAT

Object: To study the effects of seed rate and row spacing on the yields of winter wheat. The effects of different periods of pre-cropping with winter wheat are also studied - Saxmundham, Oldershaw's and Garner's plots.

The sixth year on this site, first year of row spacing and seed rate treatments, winter wheat.

For previous years see 66/C/30(t), 67/C/23(t), 68/C/39 and 69-70/S/CS/1.

Design: A single replicate of 4 x 2 x 4 in 4 blocks of 4 plots each split lengthways into 2, plus one additional plot (not split into 2) per block. Additionally, all the plots are split breadthways into 3.

Whole plot dimensions: 5.49 x 40.2. Sub plot area harvested: 0.00273.

Treatments:-

Additional plots: One immediately previous continuous winter wheat crop (1), row spacing 18.9 cm (8.25 inches) and seed rate 212 kg (S), and N to sub plots as shown below, and all combinations of:-

Whole plots:

1. Cropping history: 2, 3, 4 and 5 years of immediately previous continuous winter wheat crops 1966-1970 (2, 3, 4, 5).

Half plots:

2. Row spacing: 15.2 cm (6 inches) (C), 30.5 cm (12 inches) (W).
3. Seed rates: 70.6 (R1), 141 (R2), 212 (R3), 283 (R4) kg.

One third plots:

4. N rates: 50.2 (N1), 100.4 (N2), 150.6 (N3) kg N as 'Nitro-Chalk'.

Basal applications: 1255 kg (0:20:20) applied to stubble in autumn before ploughing. Weedkiller: Metoxuron ('Dosanex' at 5.6 kg in 449 l).

Cultivations, etc.: Basal PK applied and ploughed in: 14 Sept, 1970. Seed drilled and basal NPK applied: 5 Oct. Weedkiller applied: 15 Apr, 1971. Test N applied: 21 Apr. Combine harvested: 25 Aug. Variety: Cappelle.

71/s/CS/1

NOTE: Green crop samples for estimates of total weight, grain straw ratios and leaf areas were taken.

Standard errors per sub plot.

Grain, tonnes/hectare: 0.337 or 5.9% (6 d.f.)

SUMMARY OF RESULTS

GRAIN: TONNES/HECTARE

	R1	R2	R3	R4	N1	N2	N3	Mean
	(±0.106)				(±0.084)*			(±0.053)
C	4.67	6.11	6.20	5.87	5.09	5.99	6.06	5.71
W	5.75	5.79	5.71	5.35	4.96	5.93	6.05	5.65
					(±0.119)*			(±0.075)
				R1	4.68	5.47	5.48	5.21
				R2	5.25	6.19	6.41	5.95
				R3	5.18	6.29	6.40	5.96
				R4	4.99	5.90	5.94	5.61
Mean (±0.060)					5.02	5.96	6.06	5.68

S 5.92

General mean: 5.73

Mean D.M. %: 89.5

* For use in horizontal and interaction comparisons only