


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1971

[Full Table of Content](#)



71/R/WS/3 Dwarf Spring Wheat, Varieties, N and Ccc - S. Wheat

Rothamsted Research

Rothamsted Research (1972) *71/R/WS/3 Dwarf Spring Wheat, Varieties, N and Ccc - S. Wheat ; Yields Of The Field Experiments 1971*, pp 259 - 260 - DOI:

<https://doi.org/10.23637/ERADOC-1-97>

71/R/WS/3

SPRING WHEAT

DWARF SPRING WHEAT, VARIETIES, N AND CCC

Object: To study the effects of CCC (chlormequat) and a range of nitrogen levels on three semi-dwarf spring wheat varieties - Long Hoos IV.

Design: A single replicate of 4 x 4 x 4 in 4 blocks of 4 whole plots, split into 4 sub-plots, with split plot confounding of 3 factor interactions with blocks.

Whole plot dimensions: 2.16 x 29.6. Sub plot area harvested: 0.00087.

Treatments:

Whole plots: 1. Varieties: Benoist 257 (B), Inia (I), Kolibri (K), VR 6/57 (V).

Sub plots: All combinations of:-

2. Nitrogen: 75, 150, 225, 300 kg N as 'Nitro-chalk'.
3. Chlormequat: None, 1.12, 2.24, 3.36 kg in 337 l.

Basal applications: 336 kg (0:20:20). Weedkiller: 2,4-D at 0.56 kg plus dichlorprop at 2.24 kg in 337 l.

Cultivations, etc.: Deep-tine cultivated on 2 occasions: 10 and 12 Oct, 1970. Seed drilled at 188 kg, PK applied: 2 Apr, 1971. N applied: 16 Apr. Weedkiller applied: 11 May. Chlormequat applied: 21 May. Combine harvested: 2 Sept. Previous crops: Spring beans 1969, potatoes 1970.

NOTES: (1) Shoot heights were measured and plant numbers counted. Samples were taken just before harvest for components of yield and dry matter.

- (2) Mildew (*Erysiphe graminis*) was severe on Inia and there was considerable bird damage on all plots.

Standard error per plot estimated from unconfounded 3 factor interaction.

Grain, tonnes/hectare: Whole plot: 0.297 or 8.6% (9 d.f.)
Sub plot: 0.166 or 4.8% (15 d.f.)

71/R/WS/3

SUMMARY OF RESULTS

GRAIN: TONNES/HECTARE

	N: KG/HA				CHLORMEQUAT: KG/HA				Mean
	75	150	225	300	0	1.12	2.24	3.36	
	(1) and (2)				(1) and (2)				(±0.149)
B	3.91	4.14	4.44	4.21	3.62	4.29	4.42	4.36	4.17
I	2.21	2.30	2.51	2.63	2.27	2.54	2.40	2.45	2.41
K	2.04	2.45	2.94	3.25	2.56	2.76	2.67	2.70	2.67
V	4.30	4.65	4.64	4.64	4.26	4.59	4.60	4.77	4.56
	N: KG/HA				(±0.083)				(±0.042)
			75		3.01	3.02	3.36	3.07	3.12
			150		3.00	3.69	3.32	3.53	3.38
			225		3.40	3.91	3.56	3.66	3.63
			300		3.29	3.57	3.86	4.02	3.68
Mean (±0.042)					3.18	3.55	3.52	3.57	3.45

Mean D.M. %: 81.9

(1) (±0.165) For use in vertical and diagonal comparisons only

(2) (±0.083) For use in horizontal and interaction comparisons only