

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1971

[Full Table of Content](#)



71/R/B/8 Times of Applying Ethirimol - Barley

Rothamsted Research

Rothamsted Research (1972) *71/R/B/8 Times of Applying Ethirimol - Barley* ; Yields Of The Field Experiments 1971, pp 281 - 282 - DOI: <https://doi.org/10.23637/ERADOC-1-97>

71/R/B/8

BARLEY

TIMES OF APPLYING ETHIRIMOL

Object: To study the effects of ethirimol, applied at defined stages of mildew (*Erysiphe graminis*) epidemic and crop growth on subsequent mildew development and yield - Pastures.

Design: 4 randomised blocks of 5 plots.

Whole plot dimensions: 4.27 x 9.14. Area harvested: 0.00260.

Treatments: No systemic fungicide (O)
Seed dressing ethirimol at 1.8 kg (D)
Spray application of 0.90 kg ethirimol in 337 l
at different times:
When mildew developing on lowest leaf sheaths (S1)
At start of flag leaf emergence (S2)
At start of ear emergence (S3)

Basal applications: 377 kg (20:10:10) combine drilled. Weedkiller: Ioxynil at 0.53 kg plus mecoprop at 1.58 kg in 225 l.

Cultivations, etc.: Ploughed: 5 Nov, 1970. Seed combine drilled at 156 kg: 12 Mar, 1971. Weedkiller applied: 12 May. Ethirimol sprays applied: S1 - 12 May, S2 - 4 June, S3 - 15 June. Combine harvested: 18 Aug. Variety: Zephyr. Previous crops: Barley 1969, spring beans 1970.

NOTE: Samples were taken for assessment of mildew (*Erysiphe graminis*) and other foliar diseases.

Standard error per plot.

Grain, tonnes/hectare: 0.383 or 7.5% (12 d.f.)

71/R/B/8

SUMMARY OF RESULTS

GRAIN: TONNES/HECTARE

Q	D	S1	S2	S3	Mean
		(±0.192)			
4.69	5.61	5.21	4.90	5.23	5.13

Mean D.M. %: 85.2