

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1971

[Full Table of Content](#)



71/R/G/1 Weedkiller and Aqueous N - Grass

Rothamsted Research

Rothamsted Research (1972) *71/R/G/1 Weedkiller and Aqueous N - Grass ; Yields Of The Field Experiments 1971*, pp 341 - 345 - DOI: <https://doi.org/10.23637/ERADOC-1-97>

71/R/G/1

GRASS

WEEDKILLER AND AQUEOUS N

Object: To study the effects of a combined spray of liquid nitrogen fertiliser and a hormone weedkiller as a top dressing on new grass - Long Hoos III.

Design: 4 randomised blocks of 28 plots.

Whole plot dimensions: 2.74 x 2.13. Area harvested: 0.00022.

Treatments. (Applied per cut): All combinations of :-

1. Weedkiller (dichlorprop/MCPA): None (H0), 1.4 (H1), 2.8 (H2), 4.2 (H3) kg total a.e. (using fan jet size 00).
2. Forms of nitrogen: Solid, as 'Nitro-Chalk' 21% N applied immediately after the weedkiller (S), liquid, as urea/ammonium nitrate (26% N), mixed with the weedkiller (L) (using fan jet size 00).
3. Levels of nitrogen: 37.7 (N1), 75.3 (N2), 113.0 (N3) kg N, together with 4 additional treatments.
LLN3 H0, LLN3 H1, LLN3 H2, LLN3 H3 where fan jet size 1 was used (L1).

NOTE: The weedkiller was applied in 337 l where solid fertiliser was used. The liquid fertiliser (with or without weedkiller) was applied as a spray at 112, 225 and 337 l for rates 1, 2 and 3 respectively.

Basal applications: 750 kg (6:15:15) broadcast.

Cultivations, etc.: Spring-tine cultivated: 17 Sept, 1970. Seed drilled at 56 kg in rows 15.2 cm (6 inches) apart, basal NPK applied: 18 Sept. Grass lightly topped: 22 Apr, 1971. All treatments applied: 29 Apr. Cut: 28 May. All treatments re-applied: 9 June. Cut: 21 July. All treatments re-applied: 2 Aug. Cut: 21 Oct. Variety: S24 Perennial Ryegrass. Previous crops: Barley 1969, fallow 1970.

NOTE: Scores were made for leaf scorch. The percentage of N in the dry grass was measured for each cut.

Standard errors per plot. Dry matter, tonnes/hectare:

1st cut:	0.704 or 11.2% (69 d.f.)
2nd cut:	0.395 or 9.8% (69 d.f.)
3rd cut:	0.331 or 10.9% (69 d.f.)
Total of 3 cuts:	0.918 or 6.9% (69 d.f.)

71/R/G/1

SUMMARY OF RESULTS

1ST CUT. DRY MATTER: TONNES/HECTARE

N: KG/HA

	S	L	37.7	75.3	113.0	Mean
	(±0.203)		(±0.249)			(±0.144)
H0	6.61	6.04	6.23	6.25	6.49	6.33
H1	6.52	6.38	6.37	6.43	6.54	6.45
H2	6.46	5.95	5.84	6.14	6.64	6.21
H3	6.42	5.78	5.87	6.21	6.21	6.10
			(±0.176)			(±0.102)
		S	6.49	6.45	6.56	6.50
		L	5.67	6.07	6.38	6.04
Mean (±0.124)			6.08	6.26	6.47	6.27

LLN3 H0 6.44
 LLN3 H1 6.68
 LLN3 H2 6.62
 LLN3 H3 6.03

General mean: 6.29
 Mean D.M.%: 17.0

71/R/G/1

2ND CUT. DRY MATTER: TONNES/HECTARE

N: KG/HA

	S	L	37.7	75.3	113.0	Mean
	(±0.114)		(±0.140)			(±0.081)
H0	4.85	3.71	3.43	4.40	5.00	4.28
H1	4.53	3.52	3.01	4.15	4.92	4.03
H2	4.43	3.57	3.01	4.07	4.92	4.00
H3	4.34	3.39	2.59	4.17	4.83	3.87
			(±0.099)			(±0.057)
		S	3.56	4.62	5.43	4.54
		L	2.46	3.78	4.41	3.55
Mean (±0.070)			3.01	4.20	4.92	4.04

L1N3 H0 4.48
 L1N3 H1 4.69
 L1N3 H2 4.65
 L1N3 H3 4.54

General mean: 4.12
 Mean D.M.%: 24.1

71/R/G/1

3RD CUT. DRY MATTER: TONNES/HECTARE

N: KG/HA

	S	L	37.7	75.3	113.0	Mean
	(±0.096)		(±0.117)			(±0.068)
H0	3.50	2.74	2.25	3.28	3.83	3.12
H1	3.41	2.80	2.21	3.40	3.71	3.11
H2	3.21	2.64	2.08	3.16	3.54	2.93
H3	3.33	2.69	2.11	3.32	3.60	3.01
			(±0.083)			(±0.048)
		S	2.57	3.65	3.87	3.36
		L	1.75	2.93	3.47	2.72
Mean (±0.058)			2.16	3.29	3.67	3.04
L1N3 H0	3.53					
L1N3 H1	3.61					
L1N3 H2	3.57					
L1N3 H3	3.17					

General mean: 3.10

Mean D.M.%: 26.6

71/R/G/1

TOTAL OF 3 CUTS. DRY MATTER: TONNES/HECTARE

N: KG/HA

	S	L	37.7	75.3	113.0	Mean
	(± 0.265)		(± 0.325)			(± 0.187)
H0	14.96	12.49	11.91	13.93	15.33	13.72
H1	14.46	12.70	11.60	13.97	15.17	13.58
H2	14.10	12.17	10.93	13.37	15.10	13.14
H3	14.08	11.86	10.57	13.70	14.64	12.97
			(± 0.230)			(± 0.133)
		S	12.63	14.71	15.86	14.40
		L	9.88	12.78	14.26	12.31
Mean (± 0.162)			11.25	13.75	15.06	13.35
LLN3 H0	14.45					
LLN3 H1	14.98					
LLN3 H2	14.84					
LLN3 H3	13.74					

General mean: 13.5
Mean D.M.%: 22.6