

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1971

[Full Table of Content](#)



71/W/RN/3 Ley/ARABLE - Leys, Potatoes, Rye, Carrots, Barley

Rothamsted Research

Rothamsted Research (1972) 71/W/RN/3 Ley/ARABLE - Leys, Potatoes, Rye, Carrots, Barley ; Yields Of The Field Experiments 1971, pp 68 - 75 - DOI: <https://doi.org/10.23637/ERADOC-1-97>

71/W/RN/3

LEY/ARABLE

Object: To compare the effects on soil fertility of rotations with or without three-year leys. The effects of the cropping systems on soil-borne pathogens are also studied - Woburn Stackyard D.

The 34th year, leys, potatoes, rye, carrots, barley.

For previous years see 'Details' 1967, 68/B/2(t), 69/W/RN/3(t) and 70/W/RN/3(t).

The first test crop was changed from barley to potatoes.

Corrective K dressings (in kg K₂O) as muriate of potash applied to first test crop, potatoes.

	No FYM half plots	FYM half plots
Continuous rotations		
Ley	126	126
Sainfoin	126	126
Arable with hay	188	188
Arable	0	0
Alternating rotations (last two rotations in order)		
Arable/ley	188	439
Arable with hay/sainfoin	126	126
Ley/arable with hay	63	63
Sainfoin/arable	188	314

Treatments to first test crop potatoes:-

1. Residues of fumigant applied to potatoes 1968 (Arable and Arable with hay only) on quarter plots: None (O), 448 kg (C) chloropicrin.
2. Fresh fumigants on eighth plots: None (O), 448 kg chloropicrin plus 11.2 kg aldicarb (F).
3. Varieties, on half plots after Ley and Sainfoin and quarter plots after Arable and Arable with hay: Maris Piper (R), Pentland Crown (S).

Treatments to rye:-

1. Residues of fumigant applied to potatoes in 1970 on quarter plots: None (O), 448 kg (F) chloropicrin.

Treatments to carrots:-

1. Residues of fumigant applied to potatoes in 1969 on quarter plots: None (O), 448 kg (C) chloropicrin.

NOTE: Red clover, S 123, replaced sainfoin after the first and second year sainfoin failed. Third year sainfoin also failed and the plots were fallowed.

71/W/RN/3

Basal manuring to both test and treatment potatoes changed to 252 kg N, 252 kg P2O5, 376 kg K2O as (13:13:20).

Cultivations, etc.:

Treatment crops.

Ley 1st year: Ploughed: 4 Sept, 1970. NPK applied: 31 Mar, 1971.

Power harrowed, seed sown at 45 kg: 22 Apr. NK applied:

11 Aug. Cut twice: 2 Aug, 8 Sept.

Ley 2nd year: NK applied: 10 Mar, 1971, 15 June, 10 Aug. Cut three times: 1 June, 2 Aug, 8 Sept.

Ley 3rd year: NK applied: 10 Mar, 1971, 15 June, 10 Aug.

Cut three times: 1 June, 2 Aug, 8 Sept.

Sainfoin 1st year: Ploughed: 4 Sept, 1970. NPK applied: 31 Mar,

1971. Power harrowed, seed drilled at 45 kg: 22 Apr. Seed

redrilled, because of bird damage, at 34 kg: 12 May. Deep-

tine cultivated because of second failure: 24 June. Red

clover sown at 45 kg: 23 July. Varieties: Sainfoin: Common,

Red clover: S 123.

Sainfoin 2nd year: Sprayed with paraquat at 0.6 kg ion in 281 l:

5 Feb, 1971. NK applied: 10 Mar. Cut once: 1 June. Deep-

tine cultivated because of crop failure: 22 July. Red clover

sown at 45 kg: 23 July. Variety: Red clover S 123.

Sainfoin 3rd year: Sprayed with paraquat at 0.6 kg ion in 281 l:

5 Feb, 1971. NK applied: 10 Mar. Sprayed with paraquat at

1.1 kg ion in 281 l: 4 June. Rotary cultivated after crop

failure: 30 June. Deep-tine cultivated: 9 Aug.

Potatoes: Ploughed: 4 Sept, 1970. NPK applied: 26 Mar, 1971.

Rotary cultivated: 29 Mar. Potatoes planted: 30 Mar.

Sprayed with linuron at 1.12 kg in 371 l: 5 May. Rotary ridged:

1 June. Sprayed with mancozeb at 1.34 kg plus demeton-s-methyl

at 245 g in 416 l: 28 June. Haulm destroyed mechanically:

23 Aug. Lifted: 24 Aug. Variety: Maris Piper.

Rye: Deep-tine cultivated twice: 29 Sept, 1970. Seed combine

drilled at 190 kg: 14 Oct. N applied 13 Apr, 1971. Seeds

hay undersown at 45 kg (AH plots): 22 Apr. Combine harvested:

31 Aug. Variety: King II.

Seeds hay: Seeds undersown in rye at 34 kg: 24 Apr, 1970. NPK

applied: 10 Mar, 1971. NK applied: 16 Aug. Cut three times:

1 June, 2 Aug, 8 Sept.

Carrots: Ploughed twice: 4 Sept, 1970, 26 Sept. NPK applied:

1 Apr, 1971. Power harrowed: 15 Apr. Seed drilled at

4.5 kg: 16 Apr. Sprayed with demeton-s-methyl at 245 g in 281 l:

19 July. Mechanically hoed: 4 - 5 June. Hand hoed twice:

7 June, 2 July. Lifted: 14 - 18 Oct. Variety: Autumn King.

71/W/RN/3

Test crops.

Potatoes, 1st test crop: Half corrective K applied to all plots except 35 and 36 after carrots: 23 Sept, 1970. Half corrective K applied to plots 35 and 36: 27 Oct. Ploughed: 3 Nov. Chloropicrin applied: 18 Dec. Remaining corrective K applied to all plots: 5 Mar, 1971. NPK applied: 26 Mar. Aldicarb applied, rotary cultivated: 29 Mar. Potatoes planted: 30 Mar. Sprayed with linuron at 1.12 kg in 371 l: 5 May. Rotary ridged: 1 June. Sprayed with mancozeb at 1.34 kg plus demeton-s-methyl at 245 g in 416 l: 28 June. Sprayed with mancozeb at 1.34 kg in 371 l: 12 Aug. Haulm destroyed mechanically: 17 Sept. Sprayed with undiluted BOV at 225 l: 20 Sept. Lifted: 28 - 30 Sept. Varieties: Maris Piper and Pentland Crown.

Barley, 2nd test crop: Magnesian limestone applied at 5020 kg: 8 Oct, 1970. Ploughed: 26 Oct. Seed combine drilled at 168 kg: 10 Mar, 1971. Sprayed with ioxynil at 0.53 kg and mecoprop at 1.58 kg in 281 l: 30 Apr. Combine harvested: 18 Aug. Variety: Julia.

NOTE: Soil samples were taken from the potato plots monthly, throughout the growing season and in the barley stubble, for counts of nematodes.

71/W/RN/3

SUMMARY OF RESULTS

RYE

GRAIN: TONNES/HECTARE

	LE	SA	AH	AR	Mean
O	4.15	3.83	3.11	3.17	3.57
D*	3.86	3.68	3.16	3.04	3.44
O	3.71	4.11	3.23	2.99	3.51
F	4.30	3.39	3.04	3.22	3.49
Mean	4.01	3.75	3.14	3.11	3.50

Mean D.M. %: 71.0

* Last applied to test crop sugar beet 1963

71/W/RN/3

POTATOES TREATMENT CROP

	LE	SA	AH	AR	Mean
TOTAL TUBERS: TONNES/HECTARE					
O	39.1	31.0	27.0	26.2	30.8
D*	43.2	34.4	30.7	30.5	34.7
Mean	41.2	32.7	28.8	28.4	32.7

% WARE: 3.81 CM (1.5 INCH) RIDDLE

O	97.8	94.7	92.8	94.4	94.9
D*	96.9	95.5	95.2	95.4	95.7
Mean	97.3	95.1	94.0	94.9	95.3

* Last applied to test crop sugar beet 1964

71/W/RN/3

POTATOES 1ST TEST CROP

TOTAL TUBERS: TONNES/HECTARE

	LE	SA	Mean	AH	AR	Mean
R	68.0	74.4	71.2	68.9	62.8	65.8
S	62.6	59.6	61.1	56.2	59.8	58.0
O	64.1	65.3	64.7	60.1	56.9	58.5
D*	66.5	68.8	67.6	65.0	65.6	65.3
O	60.1	63.0	61.5	56.0	56.1	56.1
F	70.5	71.0	70.8	69.1	66.5	67.8
O				59.4	59.3	59.3
C				65.7	63.2	64.5
Mean	65.3	67.0	66.2	62.6	61.3	61.9

* Last applied to test crop sugar beet 1966

71/W/RN/3

POTATOES 1ST TEST CROP

% WARE: 3.81 CM (1.5 INCH) RIDDLE

	LE	SA	Mean	AH	AR	Mean
R	98.0	97.6	97.8	97.6	97.4	97.5
S	97.6	97.2	97.4	96.7	97.5	97.1
O	97.8	97.6	97.7	97.0	97.2	97.1
D*	97.7	97.2	97.5	97.3	97.7	97.5
O	97.8	97.4	97.6	96.9	97.3	97.1
F	97.8	97.4	97.6	97.3	97.7	97.5
O				96.9	97.2	97.1
C				97.3	97.7	97.5
Mean	97.8	97.4	97.6	97.1	97.5	97.3

* Last applied to test crop sugar beet 1966

71/W/RN/3

CARROTS

ROOTS: TONNES/HECTARE

	O	D*	O	C	Mean
SA	97.5	97.2	92.2	102.5	97.4
AR	89.3	96.4	89.4	96.3	92.9
Mean	93.4	96.8	90.8	99.4	95.1

* Last applied to test crop sugar beet 1967