


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1972

[Full Table of Content](#)



72/R/RN/9 Cereal Disease Reference Plots - W. Wheat, S. Wheat, Oats, Beans

Rothamsted Research

Rothamsted Research (1973) *72/R/RN/9 Cereal Disease Reference Plots - W. Wheat, S. Wheat, Oats, Beans* ; Yields Of The Field Experiments 1972, pp 99 - 100 - **DOI:**
<https://doi.org/10.23637/ERADOC-1-62>

72/R/RN/9

CEREAL DISEASE REFERENCE PLOTS

Object: To study the effects of intensive cereal cropping on the incidence of soil-borne diseases, especially in relation to seasonal variation - Pennell's Piece.

Sponsors: D.B. Slope, E.W. Broom.

The tenth year, winter wheat, spring wheat, oats, beans.

For previous years see 63/C/10(t), 64-65/C/9, 66/C/7, 67-68/C/5 and 69-71/R/RN/9.

Whole plot dimensions: 17.1 x 4.27. Area harvested: Winter wheat: 0.00479, Spring wheat: 0.00473.

Seed: Winter wheat: Cappelle sown at 200 kg.
Spring wheat: Kolibri sown at 190 kg.
Oats: Manod sown at 190 kg.
Beans: Maris Bead sown at 220 kg.

Cultivations, etc.: All plots sprayed with paraquat at 0.56 kg ion in 220 1: 10 Sept, 1971. Ploughed: 22 Sept.
Winter wheat: Seed combine drilled: 8 Oct, 1971. N applied: 4 Apr, 1972. Sprayed with ioxynil at 0.63 kg and mecoprop at 1.9 kg in 220 1: 28 Apr. Combine harvested: 31 Aug.
Spring wheat: Seed combine drilled: 22 Mar, 1972. N applied: 4 Apr. Sprayed with ioxynil at 0.53 kg and mecoprop at 1.6 kg in 220 1: 19 May. Combine harvested: 5 Sept.
Oats: Seed combine drilled: 18 Mar, 1972. N applied: 4 Apr. Combine harvested: 31 Aug.
Beans: Seed placement drilled: 22-Mar, 1972. Sprayed with demeton-s-methyl at 0.25 kg in 440 1: 24 June. Combine harvested: 27 Sept.

NOTES: (1) Yields were taken for winter and spring wheat only (crop sequences 2, 3, 4 and 6).
(2) Estimates were made in spring and summer of incidence of take-all (*Gaeumannomyces graminis* formerly *Ophiobolus graminis*) and eyespot (*Cercospora herpotrichoides*).

72/R/RN/9

TABLES OF MEANS

GRAIN: TONNES/HECTARE

Crop in	CROP SEQUENCE				Mean
	2	3	4	6	
1963	W	W	BE	W	
1964	W	BE	O	W	
1965	BE	O	W	W	
1966	O	W	W	W	
1967	W	W	W	W	
1968	W	W	BE	W	
1969	W	BE	O	W	
1970	BE	O	W	W	
1971	O	W	W	W	
	WINTER WHEAT				
	5.68	5.44	5.80	5.30	5.55
	SPRING WHEAT				
	3.06	2.10	3.35	2.42	2.73

Mean D.M. %: Winter wheat: 85.6
Spring wheat: 82.5