


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1973

[Full Table of Content](#)



73/R/RN/8 - Cultivation/WEEDKILLER - Wheat

Rothamsted Research

Rothamsted Research (1974) 73/R/RN/8 - Cultivation/WEEDKILLER - Wheat ; Yields Of The Field Experiments 1973, pp 110 - 112 - DOI: <https://doi.org/10.23637/ERADOC-1-98>

73/R/RN/8

CULTIVATION/WEEDKILLER

Object: To determine the long term effects of weedkillers and different methods of primary cultivation on a rotation of crops - Great Harpenden I.

Sponsors: J.R. Moffatt, G.V. Dyke, J.A. Currie.

The 13th year, winter wheat.

For previous years see 'Details' 1967, 68/B/6(t), 69/R/RN/8(t), 70/R/RN/8, 71/R/RN/8(t), and 72/R/RN/8.

Design: 2 randomised blocks of 12 plots, split into 2.

Whole plot dimensions: 12.8 x 15.2. Sub plot area harvested: 0.00434.

Treatments: All combinations of:-

Whole plots: (1) Primary cultivations annually:- CULTIVTN

Ploughed: 18 Oct, 1972, power harrowed	
and disced: 24 Oct	Plough
Rotary cultivated: 9 Nov	Rotavate
Deep-tine cultivated twice: 18 Oct, spring-	
tine cultivated: 23 Oct	Deeptine

(2) Weed control to beans 1972:- WEEDKLLR(72)

Mechanical	Mechanc1
Simazine at 1.12 kg	Simazine
Dinoseb acetate at 6.73 kg	Dinoseb

Sub plots: (3) Paraquat to bean stubble:- PARAQUAT

None	0.00
0.56 kg ion in 220 l: 5 Oct	0.56

(4) Weed control to wheat in 1973:- WEEDKLLR(73)

None	None
Ioxynil at 0.63 kg and mecoprop at	
1.9 kg in 220 l: 17 Apr, 1973	Ioxy/mec

73/R/RN/8

together with three extra plots, all of which received simazine at 1.12 kg to beans 1972:-

Spike rotary cultivated: 17 Oct, with sub plot treatments (3) and (4) above	EXTRA
Shallow ploughed: 18 Oct, power harrowed and disced: 24 Oct, whole plot received paraquat as (3) above, sub plot treatments (4) only	Spiked
Standard cultivation, ploughed for wheat: 18 Oct, power harrowed and disced: 24 Oct, with sub plot treatments (3) and (4)	ShalloPl
	Standard

NOTE: From 1973 only one phase of the rotation, wheat, potatoes, barley, beans, is maintained each year.

Basal applications: Manures: (10:24:24) at 250 kg combine drilled. 'Nitro-Chalk' at 375 kg.

Seed: Bouquet, sown at 200 kg.

Cultivations, etc.: Seed sown: 9 Nov, 1972. N applied: 12 Apr, 1973. Combine harvested: 24 Aug.

NOTE: Scores were made for weed infestation on all sub plots.

Standard errors per plot (omitting EXTRA plots). Grain, tonnes/hectare:

Whole plot: 0.476 or 8.0% (8 d.f.)

Sub plot: 0.411 or 6.9% (8 d.f.)

73/R/RN/8

TABLES OF MEANS

GRAIN: TONNES/HECTARE

	Plough	CULTIVTN Rotavate	Deeptine	Mean	Spiked	EXTRA ShalloPl	Standard
Mean	5.48	6.33	6.10	5.97	5.36	5.06	5.80
WEEDKILR(72)							
Mechancl	5.91	6.48	6.23	6.21			
Simazine	5.61	6.03	5.70	5.78			
Dinoseb	4.92	6.50	6.38	5.93			
PARAQUAT							
0.00	5.44	6.40	5.95	5.93	5.16		5.56
0.56	5.51	6.27	6.25	6.01	5.45	5.17	5.94
WEEDKILR(73)							
None	5.29	6.09	5.70	5.70	5.22	4.62	5.64
Ioxy/mec	5.67	6.58	6.50	6.25	5.50	5.49	5.97

STANDARD ERRORS OF DIFFERENCES

CULTIVTN	WEEDKILR(72)	PARAQUAT	WEEDKILR(73)	CULTIVTN	WEEDKILR(72)	PARAQUAT	CULTIVTN	WEEDKILR(73)
----------	--------------	----------	--------------	----------	--------------	----------	----------	--------------

0.275 0.275 0.137 0.137 0.476
 Except when comparing means with same level of: CULTIVTN

Grand mean 5.83

Mean D.M. % 86.6