

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1973

[Full Table of Content](#)



73/W/RN/13 - Intensive Cereals - Ley, Potatoes, Wheat, Barley

Rothamsted Research

Rothamsted Research (1974) 73/W/RN/13 - *Intensive Cereals - Ley, Potatoes, Wheat, Barley* ; Yields Of The Field Experiments 1973, pp 128 - 134 - DOI: <https://doi.org/10.23637/ERADOC-1-98>

73/W/RN/13

INTENSIVE CEREALS

Object: To study the effects of intensive cereal cropping on yield, incidence of soil-borne diseases and organic matter in the soil - Woburn Stackyard I.

Sponsors: G.W. Cooke, D.B. Slope.

The eighth year, ley, potatoes, winter wheat, barley.

For previous years see 66/B/9(t), 67/B/9, 68/B/7(t), 69/W/RN/13(t), 70/W/RN/13(t), 71/W/RN/13(t) and 72/W/RN/13.

Design: For each experiment: 2 randomised blocks of 6 plots, split into 4.

Whole plot dimensions: 8.53 x 20.4. Sub plot area harvested:
Potatoes - 0.00139. Wheat - 0.00277. Barley - 0.00273.
Ley - 0.00089.

Treatments:-

One experiment on winter wheat on part of the site of the classical wheat experiment 1877-1954

WHEATSTE

One experiment on barley on part of the site of the classical barley experiment 1877-1954

BARLYSTE

Factors tested on both experiments are the same but crop and nitrogen rates differ. All combinations of:-

Whole plots: 1. Previous crops:

PREVCROP

1966	1967	1968	1969	1970	0971	1972	
L	P	C	C	C	L	P	C/C/L/P
P	C	C	C	L	P	C	C/L/P/C
C	C	C	L	P	C	C	L/P/C/C
C	C	L	P	C	C	C	P/C/C/C
C	L	P	C	C	C	L	C/C/C/L
C	C	C	C	C	C	C	C/C/C/C

Ley = 1 year ley P = Potatoes C = Cereal: wheat or barley.

Sub plots: 2. Nitrogen fertiliser (kg N):

N

To wheat	To barley	Wheat	Barley
63	50	63	50
126	100	126	100
189	150	189	150
252	200	252	200

Ley and potatoes received basal N only, residues of dressings to cereals are tested (NRESID).

73/W/RN/13

Basal applications: All crops: Manures: (0:14:28) at 130 kg P2O₅, 260 kg K₂O, half ploughed in, half applied to the plough furrow. Weedkiller to all crops except potatoes: Paraquat at 0.56 kg ion in 280 l.

Ley: 'Nitro-Chalk' at 60 kg N in seedbed, 60 kg N after sowing, and 60 kg N after the first cut. Two cuts were taken.

Potatoes: 'Nitro-Chalk' at 150 kg N. Weedkiller: Linuron at 1.2 kg plus paraquat at 0.42 kg ion in 280 l. Fungicide with insecticide: Mancozeb at 1.3 kg plus demeton-s-methyl at 0.25 kg in 390 l. Fungicide: Mancozeb at 1.3 kg in 390 l on the first occasion and in 370 l on the second occasion.

Wheat: Weedkiller: Ioxynil at 0.63 kg with mecoprop at 1.9 kg in 280 l.

Barley: Weedkiller: Ioxynil at 0.53 kg with mecoprop at 1.6 kg in 280 l.

Seed: Ley: Italian ryegrass S22, sown at 40 kg.

Potatoes: Majestic.

Wheat: Cappelle, sown at 200 kg.

Barley: Julia, dressed with ethirimol, sown at 160 kg.

Cultivations, etc.: All plots: Half PK applied: 4 Oct, 1972.

Ploughed: 10 Oct. Remaining PK applied: 17 Oct.

Ley: Paraquat applied: 6 Sept, 1972. Power harrowed, wheat site only: 18 Oct. N applied: 16 Mar, 1973. Seeds sown: 20 Mar. N applied: 18 May, 3 July. Cut twice: 2 July, 28 Aug.

Potatoes: Power harrowed, wheat site only: 18 Oct, 1972. N applied: 23 Mar, 1973. Rotary cultivated: 6 Apr. Potatoes planted: 7 Apr. Linuron with paraquat applied: 14 May. Grubbed: 1 June. Rotary ridged: 16 June. Fungicide with insecticide applied: 5 July. Fungicide applied: 26 July, 13 Aug. Haulm mechanically destroyed: 19 Sept. Sprayed with undiluted BOV at 220 l: 24 Sept. Lifted: 4 Oct.

Wheat: Paraquat applied: 6 Sept, 1972. Power harrowed: 18 Oct. Seed sown: 20 Oct. N applied: 12 Apr, 1973. Ioxynil with mecoprop applied: 26 Apr. Combine harvested: 22 Aug.

Barley: Paraquat applied: 6 Sept, 1972. Seed sown: 12 Mar, 1973. N applied: 16 Mar. Ioxynil with mecoprop applied: 11 May. Combine harvested: 13 Aug.

NOTE: Estimates of eyespot (*Cercospora herpotrichoides*) and take-all (*Gaeumannomyces graminis*) were made on both cereal crops in July.

Standard errors per sub plot.

Wheat, grain, tonnes/hectare: 0.141 or 3.8% (12 d.f.)

Barley, grain, tonnes/hectare: 0.256 or 5.6% (12 d.f.)

73/W/RN/13

TABLE OF MEANS

LEY

DRY MATTER: TONNES/HECTARE

PERMANENT WHEAT BLOCKS

NRESID

63	126	189	252	Mean
1ST CUT				
3.79	3.77	3.74	4.35	3.91
2ND CUT				
3.68	4.07	3.98	4.19	3.98
TOTAL OF 2 CUTS				
7.47	7.84	7.72	8.54	7.89
Mean D.M. %	1st Cut:	17.6		
	2nd Cut:	21.1		
	Total of 2 cuts:	19.4		

73/W/RN/13

LEY

DRY MATTER: TONNES/HECTARE

PERMANENT BARLEY PLOTS

NRESID

50	100	150	200	Mean
1ST CUT				
5.30	5.10	5.25	4.81	5.11
2ND CUT				
4.60	4.47	4.47	4.41	4.49
TOTAL OF 2 CUTS				
9.90	9.57	9.71	9.22	9.60
Mean D.M. %	1st Cut:	16.5		
	2nd Cut:	20.4		
	Total of 2 cuts:	18.5		

73/W/RN/13

POTATOES

PERMANENT WHEAT BLOCKS

NRESID

63	126	189	252	Mean
TOTAL TUBERS: TONNES/HECTARE				
49.3	46.5	49.7	46.4	48.0
PERCENTAGE WARE: 3.81 CM (1.5 INCH) RIDDLE				
90.9	91.7	93.2	92.2	92.0

PERMANENT BARLEY BLOCKS

NRESID

50	100	150	200	Mean
TOTAL TUBERS: TONNES/HECTARE				
47.1	47.8	51.4	47.6	48.5
PERCENTAGE WARE: 3.81 CM (1.5 INCH) RIDDLE				
92.2	92.1	91.7	91.5	91.9

73/W/RN/13

WINTER WHEAT

	N				
	63	126	189	252	Mean
PREVCROP	GRAIN: TONNES/HECTARE				
C/C/L/P	4.48	5.13	4.54	4.11	4.57
C/L/P/C	3.12	4.36	4.01	4.05	3.88
L/P/C/C	2.32	3.67	3.38	3.21	3.15
C/C/C/C	2.90	3.37	3.41	3.15	3.21
Mean	3.21	4.13	3.84	3.63	3.70

STANDARD ERRORS OF DIFFERENCES

N	PREVCROP*
N	
0.071	0.141

* Within the same level of PREVCROP only

PREVCROP	STRAW: TONNES/HECTARE				
C/C/L/P	4.50	5.04	5.05	4.83	4.85
C/L/P/C	3.47	4.85	3.21	4.62	4.04
L/P/C/C	2.72	4.09	4.45	4.53	3.95
C/C/C/C	3.57	3.59	3.86	3.93	3.74
Mean	3.56	4.39	4.14	4.48	4.14

Mean D.M. % Grain 84.4
Straw 91.2

73/W/RN/13

BARLEY

N

	50	100	150	200	Mean
PREVCROP	GRAIN: TONNES/HECTARE				
C/C/L/P	4.83	4.98	4.98	4.57	4.84
C/L/P/C	3.86	5.03	4.60	4.73	4.56
L/P/C/C	3.89	4.91	4.52	4.56	4.47
C/C/C/C	3.62	4.77	4.69	4.66	4.44
Mean	4.05	4.92	4.70	4.63	4.58

STANDARD ERRORS OF DIFFERENCES

N	PREVCROP* N
0.128	0.253

* Within the same level of PREVCROP only

PREVCROP	STRAW: TONNES/HECTARE				
C/C/L/P	3.23	4.02	4.09	4.32	3.92
C/L/P/C	2.12	3.97	3.74	4.49	3.58
L/P/C/C	2.10	4.13	3.58	4.03	3.46
C/C/C/C	2.07	3.35	3.58	3.86	3.21
Mean	2.38	3.87	3.75	4.18	3.54

Mean D.M. % Grain 84.4
Straw 90.3