


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1973

[Full Table of Content](#)



73/R/CS/13 - N Levels to Old Grass - Old Grass

Rothamsted Research

Rothamsted Research (1974) *73/R/CS/13 - N Levels to Old Grass - Old Grass* ; Yields Of The Field Experiments 1973, pp 155 - 158 - DOI: <https://doi.org/10.23637/ERADOC-1-98>

73/R/CS/13

N LEVELS TO OLD GRASS

Object: To study the effects of a range of nitrogen rates on yield and botanical composition of very old permanent pasture given a single dressing of P and K annually. N fixed by legumes is estimated and the effect of treatments on nutrients available in the soil is also studied - Park Grass Old Plot 6.

Sponsors: A.E. Johnston, R.C. Flint.

The ninth year, old grass.

For previous years see 65/C/33(t), 66/C/14, 67/C/10(t), 68/C/8(t), 69/R/CS/13(t), 70/R/CS/13(t), 71/R/CS/13, 72/R/CS/13(t).

Design: 4 randomised blocks of 10 plots.

Whole plot dimensions: 1.83 x 10.1. Area harvested: 0.00090.

Treatments: Fertiliser nitrogen (kg N-total per annum applied in four equal dressings as 'Nitro-Chalk'):

	TOTAL N
None (sprayed with mecoprop to control legumes, two plots per block)	0(S)
None (two plots per block)	0
75	75
150	150
225	225
300	300
375	375
450	450

NOTE: Mecoprop applied 18 Apl, 16 July as 'Clovotox' at 11.2 l in 340 l.

Basal applications: 34 kg P as superphosphate, 224 kg K as potassium sulphate, 11 kg Mg as magnesium sulphate.

Cultivations, etc.: Basal P K Mg applied: 8 Dec, 1972. Cut: 16 May, 1973, 27 June, 7 Aug, 9 Oct. N applied: 2 Mar and then after each cut except the last.

Standard errors per plot. Dry matter, tonnes/hectare:

1st cut:	0.221 or 9.4% (29 d.f.)
2nd cut:	0.265 or 9.9% (29 d.f.)
3rd cut:	0.196 or 9.4% (29 d.f.)
4th cut:	0.163 or 12.1% (29 d.f.)
Total of 4 cuts:	0.462 or 5.5% (29 d.f.)

73/R/CS/13

TABLES OF MEANS

DRY MATTER: TONNES/HECTARE

1ST CUT

TOTAL N

O(S)	O	75	150	225	300	375	450	Mean
0.31	1.39	1.37	2.00	2.82	4.00	4.87	5.13	2.36

STANDARD ERRORS OF DIFFERENCES

TOTAL N

O(S) v O	0.111
Between any of remainder	0.156
O(S) or O v any of remainder	0.135

2ND CUT

0.78	3.02	2.69	2.78	3.26	3.59	3.35	3.41	2.67
------	------	------	------	------	------	------	------	------

STANDARD ERRORS OF DIFFERENCES

TOTAL N

O(S) v O	0.133
Between any of remainder	0.187
O(S) or O v any of remainder	0.162

Mean D.M. %	1st cut	23.8
	2nd cut	18.0

73/R/CS/13

DRY MATTER: TONNES/HECTARE

3RD CUT

TOTAL N

O(S)	0	75	150	225	300	375	450	Mean
0.42	1.87	1.94	2.41	2.49	3.17	3.03	3.33	2.10

STANDARD ERRORS OF DIFFERENCES

TOTAL N

O(S) v 0	0.098
Between any of remainder	0.139
O(S) or 0 v any of remainder	0.120

4TH CUT

0.42	1.30	1.32	1.63	1.61	1.69	1.83	1.92	1.35
------	------	------	------	------	------	------	------	------

STANDARD ERRORS OF DIFFERENCES

TOTAL N

O(S) v 0	0.082
Between any of remainder	0.115
O(S) or 0 v any of remainder	0.100

Mean D.M. %	3rd cut	19.8
	4th cut	22.8

73/R/CS/13

DRY MATTER: TONNES/HECTARE

TOTAL OF 4 CUTS

TOTAL N								
O(S)	0	75	150	225	300	375	450	Mean
1.93	7.58	7.32	8.82	10.19	12.46	13.08	13.79	8.47

STANDARD ERRORS OF DIFFERENCES

TOTAL N	
O(S) v 0	0.231
Between any of remainder	0.327
O(S) or 0 v any of remainder	0.283

Mean D.M. % 21.1