


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1973

[Full Table of Content](#)



73/R/BS/3 - Systemic Fungicide Study - Spring Barley

Rothamsted Research

Rothamsted Research (1974) *73/R/BS/3 - Systemic Fungicide Study - Spring Barley* ; Yields Of The Field Experiments 1973, pp 324 - 325 - DOI: <https://doi.org/10.23637/ERADOC-1-98>

73/R/BS/3

SPRING BARLEY

SYSTEMIC FUNGICIDE STUDY

Object: To study the effectiveness of different methyl benzimidazol-2-ylcarbamic acid (MBC) precursors and to relate chemical measurements of persistence, movement and conversion to MBC to field performance - Great Harpenden II.

Sponsors: I.J. Graham-Bryce, I.H. Williams.

Design: 6 randomised blocks of 4 plots.

Whole plot dimensions: 2.41 x 9.14. Area harvested: 0.00151.

Treatments: Fungicidal seed dressings:	FUNGICIDE
None	O
Benomyl	B
NF 48	N
Thiophanate methyl	T

All treatments applied at 0.9 kg/126 kg seed.

Basal applications: Manures: 440 kg (20:15:15) combine drilled. Weed-killer: MCPA, mecoprop and dicamba ('Tetrallex plus' at 7.0 l in 220 l).

Seed: Sultan (smut infected seed, not dressed apart from treatments) sown at 160 kg.

Cultivations, etc.: Deep-tine cultivated: 25 Aug, 1972. Ploughed: 24 Nov. Seed sown: 17 Mar, 1973. Weedkiller applied: 15 May. Combine harvested: 11 Aug. Previous crops: Beans 1971, barley 1972.

NOTE: Plant and soil samples were taken for residue analyses of the three fungicides. Counts of mildew were made three times and of smut once.

Standard error per plot.

Grain, tonnes/hectare: 0.262 or 4.9% (15 d.f.)

73/R/BS/3

TABLES OF MEANS

GRAIN: TONNES/HECTARE

FUNGICIDE				
O	B	N	T	Mean
4.88	5.57	5.71	5.34	5.37

STANDARD ERROR OF DIFFERENCES

FUNGICIDE

0.152

Mean D.M. % 86.1