


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1973

[Full Table of Content](#)



73/W/P/3 & R/P/22 384 - Varieties - Potatoes

Rothamsted Research

Rothamsted Research (1974) *73/W/P/3 & R/P/22 384 - Varieties - Potatoes* ; Yields Of The Field Experiments 1973, pp 387 - 387 - DOI: <https://doi.org/10.23637/ERADOC-1-98>

73/R/P/4

POTATOES

SPACING AND FERTILISER

Object: To study the effects of different spacings within and between rows and of fertiliser rates on yields and size grades of two potato varieties - Stackyard.

Sponsors: G.A. Hide, J.R. Moffatt, F.V. Widdowson.

Design: 3 randomised blocks of 24 plots.

Whole plot dimensions: Narrow rows 2.84 x 15.2

Wide rows 3.66 x 15.2

Plot area harvested: Narrow rows 0.00217

Wide rows 0.00279

Treatments: All combinations of:

1. Varieties:

VARIETY

King Edward

Edward

Pentland Crown

Crown

2. Compound fertiliser (13:13:20) kg:

FERT

1510

1510

3020

3020

3. Row spacing cm:

ROWSPACE

71 (28 inches)

71

91 (36 inches)

91

4. Number of plants per hectare:

PLANT/HA

37100 (by spacing seed at 38 cm on rows 71 cm apart, 30 cm on rows 91 cm apart)

37100

29700 (seed 48 cm on 71 cm rows, 38 cm on 91 cm rows)

29700

22200 (seed 61 cm on 71 cm rows, 48 cm on 91 cm rows)

22200

Basal applications: Weedkillers: Linuron at 1.2 kg with paraquat at 0.4 kg ion in 450 l. Fungicide: Mancozeb at 1.3 kg in 430 l on narrow rows and in 340 l on wide rows. Insecticide: Demeton-s-methyl at 0.25 kg applied with the fungicide.