


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1975

[Full Table of Content](#)



75/ R/PG/5 - Park Grass - Old Grass

Rothamsted Research

Rothamsted Research (1976) 75/ R/PG/5 - Park Grass - Old Grass ; Yields Of The Field Experiments 1975, pp 26 - 29 - DOI: <https://doi.org/10.23637/ERADOC-1-141>

75/R/PG/5

PARK GRASS

Object: To study the effects of organic and inorganic manures on old grass (for hay). The effects of liming are also studied.

The 120th year, hay.

For previous years see 'Details' 1967, 68/A/6(t), 69-71/R/PG/5, 72/R/PG/5(t) and 73-74/R/PG/5.

Treatments:

Whole plots: Fertilisers and organic manures:-

		MANURE
Plot 1	N1	N1
Plot 2	None (D until 1863)	O(D)
Plot 3	None	O/PLOT3
Plot 4-1	P	P
Plot 4-2	N2 F	N2P
Plot 6	N1 P K Na Mg	N1MIN
Plot 7	P K Na Mg	MIN
Plot 8	P Na Mg	PNAMG
Plot 9	N2 P K Na Mg	N2MIN
Plot 10	N2 P Na Mg	N2FNAMG
Plot 11-1	N3 P K Na Mg	N3MIN
Plot 11-2	N3 P K Na Mg Si	N3MINSi
Plot 12	None	O/PLOT12
Plot 13	D/F	D/F
Plot 14	N2* P K Na Mg	N2*MIN
Plot 15	P K Na Mg (N2* until 1875)	MIN(N2*)
Plot 16	N1* P K Na Mg	N1*MIN
Plot 17	N1*	N1*
Plot 18	N2 K Na Mg	N2KNAMG
Plot 19	D	D
Plot 20	D/N*P K	D/N*PK

N1, N2, N3: 48, 96, 144 kg N as sulphate of ammonia

N1*, N2*: 48, 96 kg N as nitrate of soda (30 kg N to Plot 20 in years with no farmyard manure)

P: 35 kg P (15 kg P to Plot 20 in years with no farmyard manure) as single superphosphate (triple superphosphate in 1974)

K: 225 kg K (45 kg K to Plot 20 in years with no farmyard manure) as sulphate of potash

Na: 15 kg Na as sulphate of soda

Mg: 10 kg Mg as sulphate of magnesia

Si: Silicate of soda at 450 kg

D: Farmyard manure at 35 tonnes every fourth year

F: Fish meal every fourth year to supply 63 kg N

MIN: P K Na Mg

75/R/PG/5

Sub plots: Liming (none to Plot 12):-

LIME

a	Ground chalk applied as necessary to maintain pH found in 1965	A
b	Ground chalk applied as necessary to achieve pH6	B
c	Ground chalk applied as necessary to achieve pH5	C
d	None	D

Additional sub plots (Plots 18, 19 and 20 only) (tonnes CaCO₃ applied every fourth year 1920-1964):-

18-1	None	N2KNAMG0
18-2	13.5	N2KNAMG2
18-3	7.9	N2KNAMG1
19-1	None	D0
19-2	6.3	D2
19-3	1.1	D1
20-1	None	D/N*PK0
20-2	5.6	D/N*PK2
20-3	1.1	D/N*PK1

Since 1965 Plot 18-1 has been split into two for treatments 'c' and 'd' above and Plot 18-3 split into two for treatments 'a' and 'b'. The remaining sub-plots of Plots 18, 19 and 20 are treated as 'a'.

NOTE: For a fuller record of treatments see 'Details' etc.

Cultivations, etc.:- P applied: 17 Jan, 1975. K applied to Plot 20, Na and Mg to Plots 6, 7, 8, 9, 10: 3 Feb. Remaining mineral fertilisers and fish meal applied: 12 Feb. N applied: 1st dressing - 25 Apr, 2nd dressing - 21 May. Cut twice: 10 June, 3 Sept.

75/R/PG/5

TABLES OF MEANS

DRY MATTER: TONNES/HECTARE

	1ST CUT LIME					2ND CUT LIME				
	A	B	C	D	MEAN	A	B	C	D	MEAN
MAINTURE										
N1	2.11	2.36	1.77	0.55	1.70	0.62	0.57	0.43	0.18	0.45
O(D)	1.69	1.83	1.16	1.06	1.44	0.26	0.35	0.23	0.21	0.26
O/PLT3	1.59	1.67	0.72	0.91	1.22	0.26	0.44	0.12	0.16	0.24
P	1.81	2.30	1.71	1.72	1.89	0.30	0.26	0.35	0.33	0.31
N2P	2.60	2.93	2.16	2.33	2.50	0.49	0.35	0.22	0.42	0.37
N1MIN	5.21	4.76			4.99	0.86	0.89			0.87
MIN	4.86	4.94	3.34	3.15	4.07	1.42	1.85	0.95	0.53	1.19
PNAMG	2.16	2.41	2.83	2.75	2.54	0.48	0.41	0.49	0.49	0.47
N2MIN	5.32	5.27	5.82	3.98	5.10	1.59	1.60	0.47	0.63	1.07
N2PNAMG	3.43	3.43	4.06	2.58	3.37	0.30	0.37	0.34	0.34	0.34
N3MIN	5.46	5.81	6.49	3.78	5.39	1.28	1.37	0.76	0.90	1.08
N3MINSI	4.95	5.33	5.72	4.10	5.03	1.61	1.69	0.75	0.81	1.22
O/PLT12	0.82		0.55		0.69	0.29		0.16		0.23
D/F	4.94	5.02	5.03	4.51	4.88	1.34	1.52	0.95	0.65	1.11
N2*MIN	3.97	4.76	4.96	4.63	4.58	0.80	1.11	1.22	1.46	1.15
MIN(N2*)	3.94		2.30		3.12	1.65		0.71		1.18
N1*MIN	4.31	4.89	4.54	4.22	4.49	1.65	1.80	1.29	1.15	1.47
N1*	2.16	2.55	2.33	1.91	2.24	0.66	0.56	0.67	0.90	0.70
N2KNAMG0			2.06	0.82	1.44			0.29	0.20	0.24
N2KNAMG2	2.05				2.05	0.58				0.58
N2KNAMG1	1.89	2.09			1.99	0.57	0.52			0.54
D0	3.10				3.10	0.57				0.57
D2	3.78				3.78	0.97				0.97
D1	3.37				3.37	0.79				0.79
D/N*PK0	4.45				4.45	0.92				0.92
D/N*PK2	4.56				4.56	0.97				0.97
D/N*PK1	4.11				4.11	1.04				1.04
MEAN DM%					23.7					42.6

75/R/PG/5

DRY MATTER: TONNES/HECTARE

TOTAL OF 2 CUTS
LIME

	A	B	C	D	MEAN
MANURE					
N1	2.72	2.93	2.20	0.73	2.15
O(D)	1.95	2.18	1.39	1.27	1.70
O/PLCT3	1.85	2.12	0.84	1.07	1.47
P	2.11	2.57	2.07	2.04	2.20
N2P	3.09	3.28	2.37	2.75	2.87
N1MIN	6.07	5.65			5.86
MIN	6.28	6.78	4.29	3.68	5.26
PNAMG	2.64	2.82	3.32	3.24	3.00
N2MIN	6.91	6.87	6.29	4.61	6.17
N2PNAMG	3.72	3.81	4.40	2.91	3.71
N3MIN	6.75	7.18	7.25	4.68	6.46
N3MIN&I	6.57	7.01	6.47	4.92	6.24
O/PLCT12	1.12		0.71		0.91
D/F	6.28	6.55	5.97	5.17	5.99
N2*MIN	4.76	5.87	6.18	6.09	5.73
MIN(N2*)	5.59		3.01		4.30
N1*MIN	5.97	6.68	5.84	5.36	5.96
N1*	2.83	3.11	3.00	2.81	2.94
N2KNAMGO			2.35	1.02	1.68
N2KNAMG2	2.63				2.63
N2KNAMG1	2.46	2.61			2.54
DO	3.67				3.67
D2	4.75				4.75
D1	4.16				4.16
D/N*PKO	5.36				5.36
D/N*PK2	5.53				5.53
D/N*PK1	5.15				5.15
MEAN DM%					33.2