


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1975

[Full Table of Content](#)



75/ R/WW/4 - Sowing Dates & Insecticides - Winter Wheat

Rothamsted Research

Rothamsted Research (1976) 75/ R/WW/4 - *Sowing Dates & Insecticides - Winter Wheat* ; Yields Of The Field Experiments 1975, pp 293 - 294 - DOI: <https://doi.org/10.23637/ERADOC-1-141>

75/R/WW/4

WINTER AND SPRING WHEAT

SOWING DATES AND INSECTICIDES

Object: To study the effects of dates of sowing and times of applying insecticides on the incidence of cereal aphids, barley yellow dwarf virus (BYDV) and yield of winter wheat - Whittlocks.

Sponsor: R.T. Plumb.

Design: 4 randomised blocks of 12 plots.

Whole plot dimensions: 6.40 x 24.4.

Treatments: All combinations of:-

1. Dates of sowing:	SOW DATE
8 November 1974	8 NOV
5 December 1974	5 DEC
27 February 1975	27 FEB
2. Phorate granules to seedbed:	INSECTICIDE(1)
None	NONE
Phorate at 5 kg	PHORATE
3. Menazon spray:	INSECTICIDE(2)
None	NONE
Menazon (0.7 l 'Saphi-Col' in 450 l on 13 June 1975)	MENAZON

NOTE: It was intended to sow winter wheat in September, October and November, but because of poor weather, only two autumn sowings were possible and the third series of plots were sown with spring wheat on 27 February.

Seed: Winter wheat: Cappelle, sown at 200 kg.
Spring wheat: Kleiber, sown at 200 kg.

Basal applications: Manures: (10:24:24) at 250 kg, 'Nitro-Chalk' at 500 kg.

Cultivations, etc.: Ploughed: 25 Sept to 9 Oct, 1974. Spring-tine cultivated twice: 7 Nov. Phorate applied, plots rotary cultivated, seed sown: 8 Nov. Phorate applied and plots power harrowed, seed sown: 5 Dec. Phorate applied, plots power harrowed, seed sown: 27 Feb, 1975. N applied to all plots: 20 Apr. Combine harvested: November and December sowing: 14 Aug, February sowing: 28 Aug. Previous crops: Barley 1973, Oats 1974.

NOTE: Aphid counts were made on 24 Apr, 22 May, 11 June, 4, 10 and 21 July, virus counts on 26 June and 4 July. Ear weights and grain numbers prior to harvest and 1,000 grain weights were determined.

75/R/WW/4

GRAIN TONNES/HECTARE

*** TABLES OF MEANS ***

INSC TCDE(1)	NONE	PHORATE	MEAN
SOW DATE			
8 NOV	5.90	5.99	5.94
5 DEC	5.59	5.74	5.66
27 FEB	3.73	4.27	4.02
MEAN	5.09	5.33	5.21

INSC TCDE(2)	NONE	MENAZON	MEAN
SOW DATE			
8 NOV	5.98	5.91	5.94
5 DEC	5.60	5.73	5.66
27 FEB	4.01	4.03	4.02
MEAN	5.20	5.22	5.21

INSC TCDE(2)	NONE	MENAZON	MEAN
INSC TCDE(1)			
NONE	5.01	5.17	5.09
PHORATE	5.39	5.28	5.33
MEAN	5.20	5.22	5.21

INSC TCDE(1)	NONE	MENAZON	PHORATE	MENAZON
INSC TCDE(2)	NONE		NONE	
SOW DATE				
8 NOV	5.94	5.86	6.02	5.96
5 DEC	5.47	5.72	5.74	5.74
27 FEB	3.61	3.94	4.41	4.13

*** STANDARD ERRORS OF DIFFERENCES OF MEANS ***

TABLE	SOW DATE	INSC TCDE(1)	INSC TCDE(2)	SOW DATE
				INSC TCDE(1)
SED	0.145	0.113	0.113	0.205

TABLE	SOW DATE	INSC TCDE(1)	SOW DATE
	INSC TCDE(2)	INSC TCDE(2)	INSC TCDE(1)
			INSC TCDE(2)
SED	0.205	0.167	0.290

*** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION ***

STRATUM	DF	SE	CV%
BLOCK.WP	33	0.410	7.9

GRAIN MEAN DM% 87.3

PLOT AREA HARVESTED 0.00390

294