

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1976

[Full Table of Content](#)

76/W/RN/15 Rotation and Fumigation - Potatoes, Barley, Sugar Beet

Rothamsted Research

Rothamsted Research (1977) *76/W/RN/15 Rotation and Fumigation - Potatoes, Barley, Sugar Beet ; Yields Of The Field Experiments 1976*, pp 106 - 111 - DOI:

<https://doi.org/10.23637/ERADOC-1-15>

76/W/RN/15

ROTATION AND FUMIGATION

Object: To study different ways of using nematicides in a three-course rotation and to determine the effects on crop yield and incidence of pathogenic nematodes - Woburn Butt Close.

Sponsors: F.G.W. Jones, A.G. Whitehead, T.D. Williams.

The eighth year, potatoes, barley, sugar beet.

For previous years see 69/W/RN/15(t), 70/W/RN/15(t) and 71-75/W/RN/15.

Design: 3 series each of 2 blocks of 3 plots split into 7.

Whole plot dimensions: 5.33 x 31.1.

Treatments:

All phases of the rotation potatoes, barley, sugar beet are present.
Each crop tests all combinations of:-

Whole plots

1. N Nitrogen fertiliser (kg N), applied cumulatively:

POTATOES & S.BEET	BARLEY	To potatoes and sugar beet	To barley
75	38	75	38
150	75	150	75
225	113	225	113

Sub plots

2. CHEMICAL Chemicals:

0	None
A (P)	Aldicarb at 6 kg before potatoes
A (SB)	Aldicarb at 6 kg before sugar beet
A (B)	Aldicarb at 6 kg before barley
A (ALL)	Aldicarb at 6 kg before all crops
DAZ(ALL)	Dazomet at 224 kg before all crops since 1970 only
BEN(ALL)	Benomyl at 22 kg before all crops since 1974 only

NOTE: Aldicarb first used in 1976. From 1969-75 dichloropropane/dichloropropene ('D-D') was applied at 448 kg to these treatments (before appropriate crops as above).

Standard applications:

Potatoes: Mamures: (0:14:28) at 1080 kg. Weedkiller: Linuron at 1.2 kg plus paraquat at 0.42 kg ion in 280 l. Insecticide: Pirimicarb at 0.14 kg in 450 l. Fungicide with insecticide: Mancozeb at 1.3 kg with demeton-s-methyl at 0.25 kg in 450 l. Fungicide: Mancozeb at 1.3 kg in 450 l.

76/W/RN/15

Barley: Manures: (0:20:20) at 310 kg combine drilled. Weedkiller: Ioxynil at 0.52 kg with mecoprop at 1.6 kg in 280 l.
Sugar beet: Manures: Magnesian limestone at 2.5 tonnes. (0:14:28) at 1080 kg. Boron at 7.4 kg B2O3 (as 'Solubor') applied with insecticide. Insecticide: Demeton-s-methyl at 0.25 kg in 390 l.

Seed: Potatoes: Pentland Crown.

Barley: Julia, dressed with ethirimol, sown at 160 kg.

Sugar beet: Klein E, sown at 6 kg.

Cultivations, etc.:-

All Series: Dazomet applied and these plots only rotary cultivated and all plots spring-time cultivated: 11 Nov, 1975.

Potatoes: Ploughed: 4 Nov, 1975. Spring-time cultivated with crumbler:

5 Nov. Ploughed: 15 Jan, 1976. Spring-time cultivated: 3 Mar.

PK applied: 15 Mar. N applied: 23 Mar. Spring-time cultivated:

24 Mar. Aldicarb and benomyl applied, all plots rotary cultivated, potatoes planted: 1 Apr. Weedkiller applied: 5 May. Grubbed: 2 June.

Rotary ridged: 3 June. Pirimicarb applied: 18 June. Fungicide with

insecticide applied: 30 June. Fungicide applied: 30 July. Haulm

mechanically destroyed: 15 Sept. Lifted: 28 Sept.

Barley: Ploughed, spring-time cultivated with crumbler: 5 Nov, 1975.

Ploughed: 15 Jan, 1976. Aldicarb and benomyl applied, all plots

rotary cultivated: 2 Mar. N applied, spring-time cultivated, seed

sown: 3 Mar. Weedkiller applied: 29 Apr. Combine harvested: 28 July.

Sugar beet: Subsoiled: Tines 140 cm apart and 56 cm deep: 1 Sept, 1975.

Magnesian limestone applied: 5 Sept. Spring-time cultivated: 18 Sept.

Ploughed twice: 16 Oct and 15 Jan, 1976. Spring-time cultivated:

3 Mar. PK applied: 15 Mar. N applied: 23 Mar. Spring-time cultivated:

24 Mar. Aldicarb and benomyl applied, all plots rotary cultivated:

1 Apr. Spring-time cultivated with crumbler, seed sown: 5 Apr.

Tractor hoed: 13 May. Singled: 20 May. Boron and insecticide applied:

28 May. Hoed by hand: 18 June, Lifted: 18 Nov.

NOTES: (1) Soil samples were taken after harvest for eelworm counts.

(2) Sugar beet: Because of rabbit damage the yields on three plots with the following treatment combinations were not taken

N	CHEMICAL
75	A(B)
225	O
225	BEN(ALL)

Estimated values were used in the analysis.

76/W/RN/15

POTATOES

TOTAL TUBERS TONNES/HECTARE

***** TABLES OF MEANS *****

N	75	150	225	MEAN
CHEMICAL				
0	11.4	11.7	10.5	11.2
A(P)	28.6	27.2	20.4	25.4
A(SB)	23.9	23.8	20.8	22.8
A(B)	18.1	19.1	12.0	16.4
A(ALL)	31.3	30.5	27.5	29.8
DAZ(ALL)	28.4	26.3	26.3	27.0
BEN(ALL)	23.3	12.9	11.7	15.9
MEAN	23.6	21.6	18.4	21.2

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	CHEMICAL	CHEMICAL N*
SED	2.81	4.87

* WITHIN THE SAME LEVEL OF N ONLY

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP.SP	18	4.87	23.0

PERCENTAGE WARE 3.81CM (1.5 INCH) RIDDLE

***** TABLES OF MEANS *****

N	75	150	225	MEAN
CHEMICAL				
0	74.6	78.3	77.5	76.8
A(P)	91.7	87.1	84.0	87.6
A(SB)	91.0	89.1	89.1	89.7
A(B)	83.1	82.7	79.0	81.6
A(ALL)	90.8	87.6	88.6	89.0
DAZ(ALL)	90.5	89.1	89.7	89.8
BEN(ALL)	85.1	83.0	74.8	81.0
MEAN	86.7	85.3	83.2	85.1

PLOT AREA HARVESTED 0.00052

76/W/RN/15

BARLEY

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

	N	38	75	113	MEAN
CHEMICAL					
0		2.02	1.91	2.35	2.09
A(P)		1.90	2.58	2.24	2.24
A(SE)		2.12	2.70	3.03	2.62
A(B)		2.24	2.69	3.25	2.72
A(ALL)		1.68	2.24	3.36	2.43
DAZ(ALL)		2.24	1.68	1.56	1.83
BEN(ALL)		2.13	2.13	2.57	2.28
MEAN		2.05	2.27	2.62	2.31

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	CHEMICAL	CHEMICAL N*
SED	0.348	0.603

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP.SP	18	0.603	26.1

GRAIN MEAN DM% 87.3

STRAW TONNES/HECTARE

***** TABLES OF MEANS *****

	N	38	75	113	MEAN
CHEMICAL					
0		1.67	1.40	2.06	1.71
A(P)		1.59	2.16	1.57	1.77
A(SB)		1.87	2.05	2.25	2.06
A(B)		1.78	2.06	2.43	2.09
A(ALL)		1.58	1.89	2.27	1.91
DAZ(ALL)		1.95	2.19	1.50	1.88
BEN(ALL)		1.97	1.97	1.96	1.97
MEAN		1.77	1.96	2.01	1.91

STRAW MEAN DM% 91.3

PLOT AREA HARVESTED 0.00052

76/W/RN/15

SUGAR BEET

ROOTS (WASHED) TONNES/HECTARE

***** TABLES OF MEANS *****

N	75	150	225	MEAN
CHEMICAL				
O	29.8	32.4	29.7	30.6
A(P)	27.7	30.9	29.8	29.5
A(SB)	31.8	36.5	37.6	35.3
A(B)	29.3	29.8	32.4	30.5
A(ALL)	33.9	37.1	36.4	35.8
DAZ(ALL)	33.3	28.0	32.4	31.3
BEN(ALL)	26.6	25.6	32.7	28.3
MEAN	30.3	31.5	33.0	31.6

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	CHEMICAL	CHEMICAL N*
SED	1.45	2.52

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP.SP	15	2.52	8.0

SUGAR PERCENTAGE

***** TABLES OF MEANS *****

N	75	150	225	MEAN
CHEMICAL				
O	15.2	14.7	14.0	14.7
A(P)	14.6	14.8	14.3	14.6
A(SB)	15.0	14.9	14.8	14.9
A(B)	15.0	14.3	14.5	14.6
A(ALL)	14.8	14.8	14.8	14.8
DAZ(ALL)	15.0	14.3	14.5	14.6
BEN(ALL)	15.0	14.7	14.4	14.7
MEAN	14.9	14.7	14.5	14.7

76/W/RN/15

SUGAR BEET

TOTAL SUGAR TONNES/HECTARE

***** TABLES OF MEANS *****

	N	75	150	225	MEAN
CHEMICAL					
O		4.54	4.78	4.15	4.49
A(P)		4.04	4.58	4.26	4.29
A(SB)		4.77	5.43	5.58	5.26
A(B)		4.45	4.28	4.69	4.47
A(ALL)		5.00	5.47	5.38	5.29
DAZ(ALL)		5.00	4.00	4.70	4.57
BEN(ALL)		3.98	3.78	4.82	4.19
MEAN		4.54	4.62	4.80	4.65

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	CHEMICAL	CHEMICAL N*
SED	0.231	0.400

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP.SP	15	0.400	8.6
PLOT AREA HARVESTED	0.00130		