

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1976

[Full Table of Content](#)

76/W/CS/138 Control of Pcn - Potatoes

Rothamsted Research

Rothamsted Research (1977) *76/W/CS/138 Control of Pcn - Potatoes ; Yields Of The Field Experiments 1976*, pp 184 - 186 - DOI: <https://doi.org/10.23637/ERADOC-1-15>

76/W/CS/138

CONTROL OF PCN

Object: To study the fresh and residual effects of several nematicides on control of *Globodera* (formerly *Heterodera*) *rostochiensis* (PCN) and yield of potatoes - Woburn Butt Close.

Sponsor: A.G. Whitehead.

The third year, potatoes.

For previous years see 74-75/W/CS/138.

Design: 4 blocks of 7 plots (NEMRES(745) balanced over blocks).

Whole plot dimensions: 2.84 x 7.01.

Treatments: All combinations of:-

1. NEMRES(745) Residues of nematicides applied in 1974 and 75:

D Z1 1	Dichloropropene at 200 kg; dazomet at 100 kg to top 7.5 cm of soil
D Z1 2	Dichloropropene at 200 kg; dazomet at 100 kg to top 15 cm of soil
D Z2 1	Dichloropropene at 200 kg; dazomet at 200 kg to top 7.5 cm of soil
D Z2 2	Dichloropropene at 200 kg; dazomet at 200 kg to top 15 cm of soil

2. NEMACIDE(76) Nematicides in 1976:

D	Dichloropropene at 200 kg
D+01	Dichloropropene at 200 kg; oxamyl at 2.8 kg
D+03	Dichloropropene at 200 kg; oxamyl at 8.4 kg
D+04	Dichloropropene at 200 kg; oxamyl at 11.2 kg

NEMACIDE(746) plus three extra treatments (applied cumulatively 1974-76):

NONE	None
(02)02	Oxamyl at 5.6 kg
(D02)D02	Dichloropropene at 200 kg; oxamyl at 5.6 kg

NOTE: The dichloropropene was injected at 20 cm and the oxamyl was applied to the surface and rotary cultivated in to a depth of 15 cm.

Basal applications: Manures: (13:13:20) at 1850 kg. Weedkiller: Linuron at 1.2 kg plus paraquat at 0.42 kg ion in 280 l. Insecticide: Pirimicarb at 0.14 kg in 450 l. Fungicide with insecticide: Mancozeb at 1.3 kg plus demeton-s-methyl at 0.25 kg in 450 l. Fungicide: Mancozeb at 1.3 kg in 450 l. Haulm desiccant: Diquat at 0.59 kg ion in 280 l.

76/W/CS/138

Seed: Pentland Crown.

Cultivations, etc.:-

Deep-tine cultivated, dichloropropene injected, all plots spring-tine cultivated: 24 Oct, 1975. Ploughed: 19 Jan, 1976. NPK applied, spring-tine cultivated: 24 Mar. Oxamyl applied, all plots rotary cultivated, potatoes planted: 23 Apr. Weedkiller applied: 7 May. Pirimicarb applied: 18 June. Fungicide plus demeton-s-methyl applied: 30 June. Fungicide applied: 30 July. Haulm desiccant applied: 6 Oct. Haulm mechanically destroyed: 11 Oct. Lifted: 22 Oct.

NOTE: Soil samples were taken in autumn 1975 and after harvest 1976 for cyst and egg counts of *Globodera* (formerly *Heterodera*) *rostochiensis*.

76/W/CS/138

TOTAL TUBERS TONNES/HECTARE

***** TABLES OF MEANS *****

NEMACIDE(76) NEMRES(745)	D	D+01	D+03	D+04	MEAN
D Z1 1	16.4	42.0	29.7	48.9	34.3
D Z1 2	4.8	36.2	30.0	34.6	26.4
D Z2 1	23.0	26.5	35.3	34.5	29.8
D Z2 2	14.2	43.3	39.8	41.7	34.7
MEAN	14.6	37.0	33.7	39.9	31.3
NEMACIDE(746)	NONE 6.4	(02)02 25.5	(D02)D02 35.3	MEAN 22.4	

GRAND MEAN 27.5

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE NEMACIDE(746)

SED 3.08

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	6	4.36	15.9

PERCENTAGE WARE 3.81CM (1.5 INCH) RIDDLE

***** TABLES OF MEANS *****

NEMACIDE(76) NEMRES(745)	D	D+01	D+03	D+04	MEAN
D Z1 1	89.0	94.6	90.9	91.2	91.4
D Z1 2	79.0	94.1	93.0	97.4	90.9
D Z2 1	96.5	87.7	93.2	95.1	93.1
D Z2 2	85.0	99.5	92.4	94.4	92.8
MEAN	87.4	94.0	92.4	94.5	92.1
NEMACIDE(746)	NONE 74.2	(02)02 88.1	(D02)D02 91.9	MEAN 84.7	

GRAND MEAN 88.9

PLOT AREA HARVESTED 0.00100