


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1977

[Full Table of Content](#)



77/R/CS/90 Cultivations for Cereals - Wheat

Rothamsted Research

Rothamsted Research (1978) *77/R/CS/90 Cultivations for Cereals - Wheat ; Yields Of The Field Experiments 1977*, pp 159 - 160 - DOI: <https://doi.org/10.23637/ERADOC-1-29>

77/R/CS/90

CULTIVATIONS FOR CEREALS

Object: To study the engineering aspects - power requirements, rate of work, revenue and costs - of different tillage systems for continuous wheat. Effects on weeds, soil pathogens and yields are also studied - Meadow.

Sponsors: D.E. Patterson (N.I.A.E.), R. Moffitt, R.D. Prew, C.A. Edwards.

The sixth year, winter wheat.

For previous years see 72-76/R/CS/90.

Design: 3 randomised blocks of 10 plots.

Whole plot dimensions: 13.7 x 33.8.

Treatments

TILLAGE Tillage systems:-

- 1 Three passages of the tractor (three-pass system): Ploughed* 20 cm deep (8 inches): disc harrowed: drilled.
- 2 Three-pass system: Chisel-ploughed once* 15 cm deep (6 inches): disc harrowed: drilled.
- 3 Two-pass system: Ploughed* 20 cm deep: spring-tine cultivated and drilled.
- 4 Two-pass system: Ploughed* 10 cm deep: spring-tine cultivated and drilled.
- 5 Two-pass system: Sprayed with paraquat (0.56 kg ion in 220 l on 20 Oct, 1976): spring-tine cultivated and drilled.
- 6 Three-pass system: Heavy spring-tine cultivated* 15 cm deep: disc harrowed: spring-tine cultivated and drilled.
- 7 Two-pass system (stubble of 1976 crop burnt on 16 Aug): Heavy spring-tine cultivated* 15 cm deep: spring-tine cultivated and drilled.
- 8 Two-pass system: Sprayed with paraquat (0.56 kg ion in 220 l on 20 Oct): direct drilled.
- 9 Three-pass system: Rotary digger (N.I.A.E.) cultivated*: disc harrowed: spring-tine cultivated and drilled.
- 10 Two-pass system (stubble of 1976 crop burnt on 16 Aug): Rotary digger (N.I.A.E.) cultivated*: spring-tine cultivated and drilled.

NOTE: Rotary digger (N.I.A.E.) - depth of working: rotor 10 cm, tines 20 cm.

* Cultivations done on 30 Sept, 1976. Three plots of TILLAGE 1, two of TILLAGE 2 and one each of TILLAGE 3, 6 and 10 received remaining cultivations and were sown on 21 Oct. TILLAGE 8 plots sown on 2 Nov. All remaining cultivations and drilling done on 3 Nov, and these plots only disc harrowed. A disc drill was used on all treatments except 8.

Basal applications: Manures: Chalk at 7.5 t. (10:24:24) at 250 kg, combine drilled. 'Nitro-Chalk' at 400 kg. Weedkillers: Paraquat at 0.56 kg ion in 220 l, methabenzthiazuron at 3.1 kg in 220 l. Insecticide: Pirimicarb at 0.14 kg in 270 l.

Seed: Maris Huntsman, sown at 160 kg.

Cultivations, etc.: Chalk applied: 1 Sept, 1976. Paraquat applied: 27 Sept. Methabenzthiazuron applied: 19 Nov. N applied: 8 Apr, 1977. Insecticide applied: 13 July. Combine harvested: 7 Sept.

77/R/CS/90

NOTES: Observations and determinations were made as follows:-

- (1) Soil: Structure assessments, photographs for aggregate size, moisture determinations.
- (2) Implements: Depth and width of work, forward speed, wheel slip and power requirements.
- (3) Crop: Plant and tiller counts, number of ears and grains per ear, disease and pest assessments, aerial photographs.

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

TILLAGE	1	2	3	4	5	6	7	8	9	10	MEAN
	5.24	5.04	5.18	5.35	4.74	4.79	5.16	4.49	5.02	5.27	5.03

***** STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	TILLAGE
-----	-----
SED	0.197

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
BLOCK.WP	18	0.242	4.8

GRAIN MEAN DM% 81.0

PLOT AREA HARVESTED 0.01031