


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1977

[Full Table of Content](#)



77/S/WW/1 Rates and Times of N and Fungicide - W. Wheat

Rothamsted Research

Rothamsted Research (1978) *77/S/WW/1 Rates and Times of N and Fungicide - W. Wheat* ; Yields Of The Field Experiments 1977, pp 329 - 331 - DOI: <https://doi.org/10.23637/ERADOC-1-29>

77/S/WW/1

WINTER WHEAT

RATES AND TIMES OF N AND FUNGICIDE

Object: To study the effects of fungicide and rates and times of applying nitrogen fertiliser on the incidence of foliar diseases and on yield - Saxmundham.

Sponsors: F.V. Widdowson, A. Penny.

Design: Single replicate of 4 x 2 x 2 x 2 fully randomised plus 4 extra plots.

Whole plot dimensions: 2.74 x 6.10.

Treatments: All combinations of:-

1. S N RATE Rates of solid nitrogen fertiliser (kg N):

0
50
100
150

2. S N TIME Times of applying solid nitrogen fertiliser:

APRIL 28 April
MAY 19 May

3. L N RATE Rates of applying liquid nitrogen fertiliser, half on 16 June, half on 6 July (kg N):

0
50

4. FUNGICIDE Foliar spray to control fungi:

NONE None
B+MB+MZ Benomyl + maneb + mancozeb

plus four extra plots not given liquid fertiliser but given benomyl + maneb + mancozeb as above, and testing solid nitrogen as follows (kg N):-

XTRA S N	G.S.3(16 Mar)	G.S.5(28 Apr)	G.S.8(25 May)			
20+60+20	20	+	60	+	20	(duplicated)
30+90+30	30	+	90	+	30	(duplicated)

NOTE: Benomyl applied at 0.28 kg with mancozeb plus maneb ('Kascade' at 2.24 kg) in 340 l on 19 May, 16 June, 6 July.

Basal applications: Manures: K20 at 150 kg as muriate of potash. (30:13:0) at 190 kg. Weedkillers: Ioxynil at 0.53 kg with mecoprop at 1.6 kg in 340 l. Growth regulator: Chlormequat at 1.7 kg in 340 l.

Seed: Maris Huntsman, broadcast at 200 kg.

Cultivations, etc.: K applied: 17 Aug, 1976. Ploughed 3 Sept. Seed broadcast, NP applied: 23 Nov. Spring-tine cultivated: 24 Nov. Weedkillers applied: 14 May, 1977. Growth regulator applied: 19 May. Combine harvested: 1 Sept. Previous crops: Beans 1975, wheat 1976.

77/S/WW/1

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

INCLUDING S N RATE 0

S N RATE	0	50	100	150	MEAN
L N RATE					
0	3.20	4.72	5.29	5.76	4.74
50	3.61	4.64	4.77	5.71	4.68
MEAN	3.40	4.68	5.03	5.74	4.71

S N RATE	0	50	100	150	MEAN
FUNGICIDE					
NONE	3.38	4.80	4.78	5.43	4.60
B+MB+MZ	3.42	4.56	5.28	6.04	4.83
MEAN	3.40	4.68	5.03	5.74	4.71

FUNGICIDE	NONE	B+MB+MZ	MEAN
L N RATE			
0	4.65	4.84	4.74
50	4.55	4.82	4.68
MEAN	4.60	4.83	4.71

EXCLUDING S N RATE 0

S N RATE	50	100	150	MEAN
S N TIME				
APRIL	5.49	5.84	6.53	5.95
MAY	3.88	4.22	4.94	4.35
MEAN	4.68	5.03	5.74	5.15

L N RATE	0	50	MEAN
S N TIME			
APRIL	6.06	5.85	5.95
MAY	4.46	4.24	4.35
MEAN	5.26	5.04	5.15

FUNGTIME	NONE	B+MB+MZ	MEAN
S N TIME			
APRIL	5.63	6.27	5.95
MAY	4.38	6.38	4.35
MEAN	5.01	5.29	5.15

XTRA S N	20+60+20	30+90+30	MEAN
	6.19	6.69	6.44

GRAND MEAN 4.91

77/S/WW/1

GRAIN TONNES/HECTARE

*****STANDARD ERRORS OF DIFFERENCES OF MEANS *****

TABLE	XTRA S N	L N RATE	FUNGCIDE	S N RATE
SED	0.369	0.131 0.151*	0.131 0.151*	0.185

TABLE	S N TIME	L N RATE FUNGCIDE	S N RATE S N TIME	S N RATE L N RATE
SED	0.151	0.185	0.261	0.261

TABLE	S N TIME L N RATE	S N RATE FUNGCIDE	S N TIME FUNGCIDE
SED	0.213	0.261	0.213

* USE ONLY WITH TABLES EXCLUDING S N RATE 0

***** STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION *****

STRATUM	DF	SE	CV%
WP	15	0.369	7.5

GRAIN MEAN DM% 76.8

PLOT AREA HARVESTED 0.00089