

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1977

[Full Table of Content](#)



77/S/RN/2 Rotation II - Wheat, Barley

Rothamsted Research

Rothamsted Research (1978) *77/S/RN/2 Rotation II - Wheat, Barley* ; Yields Of The Field Experiments 1977, pp 41 - 45 - DOI: <https://doi.org/10.23637/ERADOC-1-29>

77/S/RN/2

ROTATION II

Object: To measure, by crop yields and soil analysis, the residual value of P applied as FYM or superphosphate in the periods 1899-1964 and 1965-1967 - Saxmundham.

Sponsors: G.E.G. Mattingly, A.E. Johnston.

The ninth year of revised scheme, wheat, barley.

For previous years see 'Details' 1967 & 1973, and 74-76/S/RN/2.

Whole plot dimensions: 5.49 x 39.8.

Treatments: From 1899-1964 the experiment tested farmyard manure and nitrogen and phosphate fertilisers applied to a rotation of crops. Since 1965 the treatments have been changed to evaluate old residues of P (from FYM and superphosphate) and new residues from treatments applied 1965-1967. All crops of the rotation - potatoes, barley, sugar beet, barley - were grown until 1974. The whole experiment was sown to barley in 1975 and 1976, wheat and barley in 1977, and tests combinations of:

Whole plots

1. RESIDUE Residues of previous treatments:-

		Approximate total dressing 1899-1964	Total dressing 1965-1967
(O)O	Plot 1	None	None
(D)O	Plot 2	400 tonnes FYM	None
(DP)O	Plot 3	400 tonnes FYM, 2.7 tonnes P205	None
(DP)D2	Plot 4	400 tonnes FYM, 2.7 tonnes P205	100 tonnes FYM
(DP)D2P1	Plot 5	400 tonnes FYM, 2.7 tonnes P205	100 tonnes FYM, 0.56 tonnes P205
(DP)P1	Plot 6	400 tonnes FYM, 2.7 tonnes P205	0.56 tonnes P205
(DP)P2	Plot 7	400 tonnes FYM, 2.7 tonnes P205	1.13 tonnes P205
(DP52)O	Plot 8	326 tonnes FYM, 4.3 tonnes P205 (until 1952 only)	None

3rd barley after potatoes or sugar beet (1974) tests in addition to 1:-

Sub plots

2. P205 72 Phosphate in 1970-72 (total P205 applied (kg)):

0	0 (2 sub plots/plot)
126	126
252	252
378	378

and some of the combinations of 2 with:-

77/S/RN/2

3. P205 746 Phosphate in 1974-76 (kg P205):

	1974	1975-6
0	None	None
63x3	63	63
189	189	None

Wheat (Maris Huntsman) after barley (1974-76) after potatoes (1973) and:
Wheat (Cappelle) after barley (1974-76) after sugar beet (1973), test in
addition to 1:

Sub plots

2. P205 71 Phosphate residues 1969-71 (total P205 applied (kg)):

	0 (2 sub plots/plot)
0	126
126	252
252	378
378	

and some of the combinations of 2 with:-

3. P205 735 Phosphate in 1973-75 (kg P205):

	1973	1974, 75
0	None	None
63x3	63	63
189	189	None

Standard applications:

Winter wheat: Manures: K20 at 150 kg as muriate of potash. N at 50 kg at
drilling as 'Nitro-Chalk'. N at 125 kg top-dressed in spring as 'Nitro-Chalk'.
Weedkillers: Ioxynil at 0.53 kg with mecoprop at 1.6 kg in 340 l.

Barley: Manures: K20 at 150 kg as muriate of potash. After barley (1975-76)
after potatoes (1974): N at 94 kg as 'Nitro-Chalk' (N2). After barley
(1975-76) after sugar beet (1974): N at 63 kg as 'Nitro-Chalk' (N1).

Weedkillers: Ioxynil at 0.42 kg with mecoprop at 1.3 kg in 340 l. Fungicide:
Tridemorph at 0.53 kg applied with the weedkillers.

Seed: Winter wheat varieties sown at 200 kg.

Barley: Julia, dressed ethirimol, sown at 190 kg.

Cultivations, etc.: K applied: 17 Aug, 1976. Ploughed: 6 Sept.

Winter wheat: Sown: 23 Nov. Spring N applied: 28 Apr, 1977. Weedkillers applied:
14 May. Harvested: 1 Sept.

Barley: Sown, N applied: 6 Apr. Weedkillers plus fungicide applied: 25 May.
Harvested: 31 Aug.

77/S/RN/2

WINTER WHEAT (MARIS HUNTSMAN) AFTER BARLEY 1974-76 POTATOES 1973

***** TABLES OF MEANS *****

		GRAIN TONNES/HECTARE			STRAW TONNES/HECTARE		
	P205 735	0	63X3	189	0	63X3	189
RESIDUE	P205 71						
(O)O	0	4.08	4.86		2.55	3.10	
(O)O	126			4.51			2.05
(O)O	252		5.72			3.81	
(O)O	378			5.11			2.57
(D)O	0	5.00		5.00	3.15		2.34
(D)O	126		6.33			3.43	
(D)O	252			5.47			2.59
(D)O	378		5.72			3.27	
(DP)O	0	6.06		6.35	3.29		3.78
(DP)O	126		6.81			4.60	
(DP)O	252			6.43			3.27
(DP)O	378		5.94			3.52	
(DP)D2	0	6.45	6.77		3.66	4.33	
(DP)D2	126			5.61			2.86
(DP)D2	252		6.91			3.77	
(DP)D2	378			6.43			3.66
(DP)D2P1	0	6.42		6.82	3.47		3.80
(DP)D2P1	126		5.97			3.52	
(DP)D2P1	252			6.69			4.06
(DP)D2P1	378		6.07			3.39	
(DP)P1	0	6.42		7.18	4.33		4.08
(DP)P1	126		6.33			3.81	
(DP)P1	252			6.67			4.29
(DP)P1	378		6.16			3.52	
(DP)P2	0	6.18	7.05		3.77	4.69	
(DP)P2	126			6.42			3.43
(DP)P2	252		7.03			4.00	
(DP)P2	378			6.67			3.75
(DP52)O	0	6.65	7.27		4.14	4.82	
(DP52)O	126			5.47			3.21
(DP52)O	252		6.42			3.79	
(DP52)O	378			6.52			4.70

GRAIN MEAN DM% 77.6

STRAW MEAN DM% 88.2

PLOT AREA HARVESTED 0.00077

77/S/RN/2

WINTER WHEAT (CAPPELLE) AFTER BARLEY 1974-6 SUGAR BEET 1973

***** TABLES OF MEANS *****

		GRAIN TONNES/HECTARE			STRAW TONNES/HECTARE		
	P205 735	0	63X3	189	0	63X3	189
RESIDUE	P205 71						
(O)O	0	2.86		3.82	2.33		2.98
(O)O	126		5.15			3.80	
(O)O	252			4.67			3.10
(O)O	378		5.02			3.63	
(D)O	0	3.68	4.30		2.88	2.83	
(D)O	126			4.51			3.12
(D)O	252		5.11			3.50	
(D)O	378			5.02			3.51
(DP)O	0	5.14	5.76		4.28	4.03	
(DP)O	126			5.54			4.06
(DP)O	252		5.66			3.30	
(DP)O	378			5.88			4.19
(DP)D2	0	6.21		6.12	3.81		4.66
(DP)D2	126		6.74			4.61	
(DP)D2	252			6.47			4.42
(DP)D2	378		6.36			4.48	
(DP)D2P1	0	6.86	6.35		4.81	4.85	
(DP)D2P1	126			6.31			4.85
(DP)D2P1	252		6.89			4.82	
(DP)D2P1	378			6.35			4.02
(DP)P1	0	6.27	6.33		4.83	4.56	
(DP)P1	126			6.50			4.40
(DP)P1	252		6.50			4.28	
(DP)P1	378			6.33			4.70
(DP)P2	0	5.87		6.24	4.16		4.19
(DP)P2	126		6.11			4.04	
(DP)P2	252			6.26			4.36
(DP)P2	378		6.82			4.75	
(DP52)O	0	6.47		6.02	4.42		3.97
(DP52)O	126		5.87			4.82	
(DP52)O	252			6.51			5.33
(DP52)O	378		5.42			4.16	

GRAIN MEAN DM% 78.1

STRAW MEAN DM% 85.6

PLOT AREA HARVESTED 0.00077

77/S/RN/2

BARLEY AFTER BARLEY 1975-6 POTATOES 1974
AND
BARLEY AFTER BARLEY 1975-6 SUGAR BEET 1974

GRAIN TONNES/HECTARE

***** TABLES OF MEANS *****

PREVIOUS CROP 1974		POTATOES(N2)				SUGAR BEET(N1)	
	P205 746	0	63X3	189	0	63X3	189
RESIDUE	P205 72						
(O)O	0	4.43		4.19	4.16	4.68	
(O)O	126		5.53				4.20
(O)O	252			4.55		4.80	
(O)O	373		4.77				4.54
(D)O	0	4.43	5.76		4.94		5.17
(D)O	126			4.65		5.42	
(D)O	252		5.00				5.27
(D)O	373			5.96		5.29	
(DP)O	0	5.13	5.23		5.06		5.27
(DP)O	126			5.61		5.79	
(DP)O	252		6.58				5.51
(DP)O	373			5.16		5.29	
(DP)D2	0	5.84		5.62	5.48	5.39	
(DP)D2	126		5.63				5.54
(DP)D2	252			6.48		5.79	
(DP)D2	373		5.85				5.55
(DP)D2P1	0	5.91	5.79		5.77		5.62
(DP)D2P1	126			5.91		5.68	
(DP)D2P1	252		6.20				6.28
(DP)D2P1	373			6.55		5.68	
(DP)P1	0	5.85	5.55		5.68		5.66
(DP)P1	126			5.03		5.79	
(DP)P1	252		6.55				5.94
(DP)P1	373			6.02		5.89	
(DP)P2	0	6.34		5.86	5.80	5.29	
(DP)P2	126		6.03				5.27
(DP)P2	252			5.89		5.91	
(DP)P2	373		5.55				5.91
(DP52)O	0	5.21		5.59	5.01	5.55	
(DP52)O	126		5.62				5.07
(DP52)O	252			5.71		5.28	
(DP52)O	373		6.11				5.54

GRAIN MEAN DM% (PREVIOUS CROP 1974 POTATOES) 79.7

GRAIN MEAN DM% (PREVIOUS CROP 1974 SUGAR BEET) 80.1

PLOT AREA HARVESTED 0.00077